

University of Toledo Medical Center Department of Orthopaedic Surgery

The University of Toledo is accredited by the Accreditation Council for Continuing Medical Education (ACCME) to provide continuing medical education for physicians. The University of Toledo designates this enduring activity for a maximum of 2 AMA PRA *Category 1 Credits*^{IM}. Physicians should claim only credit commensurate with the extent of their participation in the activity. *Termination Date for CME July 30, 2019*

Name:	Degree:
Specialty:	
Address:	
Phone (Required):	Email (Required):

Learning Objectives: Participants will be able to:

- Describe common orthopaedic concerns
- Review new treatment options available for common orthopaedic injuries.

OCTOBER 2016 Orthopaedic Monthly Questions

Please check the appropriate answer and mail back to the CME office to receive credit. You must obtain at least 80% (4/5) to receive CME credit. There is a \$10.00 processing fee for CME credit.

- 1. Fracture of the femoral neck may cause
 - A) Vascular injury.
 - B) Osteonecrosis.
 - C) Compartment syndrome.
 - D) Progressive valgus deformity.
- 2. What are some associated conditions of scapular fractures?
 - A) Pneumothorax
 - B) Head injury
 - C) Pulmonary contusion
 - D) All of the Above
- 3. What is another name for Floating Shoulder?
 - A) Injury of the Superior Shoulder Suspensory Complex
 - B) Scapulothoracic Dissociation
 - C) Coracoid Fracture
 - D) None of the Above

To receive CME Credit complete this post-test and pay via:

Check: mail completed post-test with \$10 payment to: Center for Continuing Medical Education, The University of Toledo, 3000 Arlington Avenue MS 1092, Toledo, OH 43614

Credit Card: email completed post test to: <u>ContinuingMedEd@utoledo.edu</u> (Include phone # so we can call for payment)

CME Credit will not be awarded until payment has been received. Once your credit has been awarded, you will receive an email from the CME Office notifying you how to obtain your credit.

- 4. The radial nerve gives innervation to the
 - A) Finger Extensors
 - B) Wrist Extensors
 - C) Thumb Extensor
 - D) All of the Above
- 5. What is the death rate associated with flail chest injuries?
 - A) 0% 10%
 - B) 10% 25%
 - C) 25% 50%
 - D) Over 50%