

THE UNIVERSITY OF TOLEDO

MEDICAL CENTER

University of Toledo Medical Center Department of Orthopaedic Surgery

The University of Toledo is accredited by the ACCME to provide continuing medical education for physicians. The University of Toledo designates this enduring activity for a maximum of **2** AMA PRA Category 1 CreditsTM. Physicians should claim only credit commensurate with the extent of their participation in the activity.

Termination Date for CME July 30, 2019

Name:	Degree:
Specialty:	
Address:	
Phone (Required):	Email (Required):
	Email (Required):

Learning Objectives: Participants will be able to:

- Describe common orthopaedic concerns
- Review new treatment options available for common orthopaedic injuries.

June 2018 Orthopaedic Monthly Questions

Please check the appropriate answer and mail back to the CME office to receive credit. You must obtain at least 80% (4/5) to receive CME credit. There is a \$10.00 processing fee for CME credit.

- 1. A scenario that would aggravate the rhomboid trigger points
 - is____
 - A) Painting
 - B) Bad Posture
 - C) Sitting in a protracted position
 - D) All of the Above
- 2. The normal Atlanto-dens interval measures between
 - A) 10-15mm
 - B) Less than 5mm
 - C) 6-8mm
 - D) 5-10mm
- 3. A ganglion cyst of the foot can be differentiated from plantar fibromatosis by:
 - A) Transillumination
 - B) The location of the mass/pain
 - C) Aspiration
 - D) All of the Above

To receive CME Credit complete this post-test and pay via:

Check: mail completed post-test with \$10 payment to: Center for Continuing Medical Education, The University of Toledo, 3000 Arlington Avenue MS 1092, Toledo, OH 43614

Credit Card: email completed post test to: <u>ContinuingMedEd@utoledo.edu</u> (Include phone # so we can call for payment)

CME Credit will not be awarded until payment has been received. Once your credit has been awarded, you will receive an email from the CME Office notifying you how to obtain your credit.

- 4. The short head of the biceps femoris is innervated by the:
 - A) Common peroneal nerve
 - B) Ulnar Nerve
 - C) Pudental Nerve
 - D) Median Nerve
- 5. The rhomboid muscle is innervated by:
 - A) The radial nerve
 - B) The ulnar nerve
 - C) The dorsal scapular nerve
 - D) The medial nerve