



Original Contribution

Involvement of Na^+/K^+ -ATPase in hydrogen peroxide-induced hypertrophy in cardiac myocytes

Lijun Liu ^{a,1}, Jie Li ^{a,1}, Jiang Liu ^b, Zhaokai Yuan ^a, Sandrine V. Pierre ^a, Weikai Qu ^a,
Xiaochen Zhao ^a, Zijian Xie ^{a,b,*}

^a Department of Physiology, Pharmacology, Metabolism and Cardiovascular Sciences, Medical University of Ohio, Toledo, OH 43614, USA

^b Department of Medicine, Medical University of Ohio, 3035 Arlington Avenue, Toledo, OH 43614, USA

Received 14 April 2006; revised 8 August 2006; accepted 21 August 2006

Available online 25 August 2006

Abstract

We have shown that increased production of reactive oxygen species (ROS) was required for ouabain-induced hypertrophy in cultured cardiac myocytes. In the present study we assessed whether long-term exposure of myocytes to nontoxic ROS stress alone is sufficient to induce hypertrophy. A moderate amount of H_2O_2 was continuously generated in culture media by glucose oxidase. This resulted in a steady increase in intracellular ROS in cultured cardiac myocytes for at least 12 h. Such sustained, but not transient, increase in intracellular ROS at a level comparable to that induced by ouabain was sufficient to stimulate protein synthesis, increase cell size, and change the expression of several hypertrophic marker genes. Like ouabain, glucose oxidase increased intracellular Ca^{2+} and activated extracellular signal-regulated kinases 1 and 2 (ERK1/2). These effects of glucose oxidase were additive to ouabain-induced cellular changes. Furthermore, glucose oxidase stimulated endocytosis of the plasma membrane Na^+/K^+ -ATPase, resulting in significant inhibition of sodium pump activity. While inhibition of ERK1/2 abolished glucose oxidase-induced increases in protein synthesis, chelating intracellular Ca^{2+} by BAPTA-AM showed no effect. These results, taken together with our prior observations, suggest that ROS may cross talk with Na^+/K^+ -ATPase, leading to the activation of hypertrophic pathways in cardiac myocytes.

© 2006 Elsevier Inc. All rights reserved.

Keywords: ROS; Na^+/K^+ -ATPase; Ouabain; Cardiac myocytes; Hypertrophy

Introduction

Na^+/K^+ -ATPase, or sodium pump, is a ubiquitous transmembrane enzyme that has long been ascribed the function of transporting Na^+ and K^+ across the plasma membrane [1,2]. Recently, we have demonstrated that Na^+/K^+ -ATPase is an important signal transducer in cardiac myocytes [3–6]. Binding of ouabain to the Na^+/K^+ -ATPase stimulates multiple pathways

including activation of phospholipase C (PLC)/protein kinase C (PKC) cascade [7] and Src/EGFR/Ras/Raf/ERK1/2 cascade [8] and increased production of ROS [9]. ROS, such as the superoxide radical and H_2O_2 , are continuously produced in most cells, and their levels are regulated by a number of enzymes and physiological antioxidants. It has been known that excessive generation of ROS is associated with cell injury in a variety of cardiovascular disorders [10–13], including atherosclerosis, ischemic heart disease, hypertension, and cardiac failure. Recently, it has become clear that ROS also act as an important second messenger in several signal pathways that control gene transcription [14,15] and cell growth [10,16–19]. ROS generation is linked to the pathways of many hypertrophic stimuli including the Na^+/K^+ -ATPase ligands [9,20,21], agonists for G_q/G_{11} -coupled receptors [22–24], growth factors [25,26], and mechanotransduction [27].

We observed that partial inhibition of Na^+/K^+ -ATPase by ouabain increased ROS production in cardiac myocytes.

Abbreviations: CM-DCFH, chloromethyl dihydrodichlorofluorescein diacetate; ERK, extracellular signal-regulated kinase; NAC, *N*-acetylcysteine; PKC, protein kinase C; PLC, phospholipase C; ROS, reactive oxygen species; SDS-PAGE, sodium dodecyl sulfate-polyacrylamide gel electrophoresis.

* Corresponding author. Department of Physiology, Pharmacology, Metabolism and Cardiovascular Sciences, Medical University of Ohio, Toledo, OH 43614, USA. Fax: +1 419 383 2871.

E-mail address: zxie@meduohio.edu (Z. Xie).

¹ These authors contributed equally to this work.

Addition of antioxidant *N*-acetylcysteine (NAC) abolished ouabain-induced protein synthesis and expression of skACT and ANF as well as ouabain-induced down-regulation of the Na⁺/K⁺-ATPase α_3 gene [9]. These data, together with other reports [26–28], indicate that ROS are involved in the activation of hypertrophic pathways. However, whether increases in ROS stress alone are sufficient to induce hypertrophic growth remains to be established. In this study, we developed a chronic ROS stress model, assessed whether a sustained and moderate increase in intracellular ROS alone can stimulate hypertrophic growth, and explored the molecular mechanism of this ROS-induced cell growth in cultured neonatal rat cardiac myocytes.

Experimental procedures

Materials

TRI reagent for RNA isolation was purchased from Molecular Research Center, Inc. (Cincinnati, OH). Radio-nucleotides (³²P labeled, about 3000 Ci/mmol) and ⁸⁶Rb⁺ were obtained from Dupont NEN (Boston, MA). [³H]Phenylalanine was purchased from Amersham (Cleveland, OH). Fura-2 AM and chloromethyl dihydrodichlorofluorescein diacetate (CM-DCFH) were obtained from Molecular Probes (Eugene, OR). The antibodies used and their sources were as follows: the polyclonal anti-ERK1/2 antibody, the monoclonal anti-phospho-ERK1/2 antibody, the goat anti-rabbit, and the goat anti-mouse-conjugated HRP secondary antibodies were obtained from Santa Cruz Biotechnology (Santa Cruz, CA). Chemiluminescence ECL kit was purchased from Pierce (Rockford, IL). The Optitran nitrocellulose membranes were obtained from Schleicher and Schuell (Keene, NH). All research on rats was done according to procedures and guidelines approved by the Institutional Animal Care and Use Committee.

Cell preparation and culture

Cultures of neonatal rat cardiac myocytes were prepared as described in our previous work [5]. Briefly, myocytes were isolated from ventricles of 1-to 2-day-old Sprague-Dawley rats and purified by centrifugation on Percoll gradients. Myocytes were seeded at a density of 1×10^5 cells/cm² and cultured in a medium containing 4 parts of DMEM and 1 part of Medium 199, penicillin (100 units/ml), streptomycin (100 μ g/ml), and 10% fetal bovine serum for 24 h, and then incubated in serum-free medium for 48 h before use for the experiments. These cultures contain more than 95% myocytes as assessed by immunofluorescence staining with a myosin heavy chain antibody.

Fluorescence microscopic measurements of intracellular Ca²⁺ concentration and ROS production

Myocytes were cultured on laminin-coated glass coverslips in 6-well plates. Intracellular Ca²⁺ was measured by fura-2 AM and intracellular ROS were assessed with the ROS-sensitive

fluorophore CM-DCFH as described previously [9]. Briefly, fura-2 AM fluorescence was recorded using an Attofluor imaging system (Atto Instruments) at excitation wavelength of 340/380 nm and at emission wavelength of 505 nm. Under each experimental condition time-averaged signals were obtained from about 40 single cells. Relative Ca²⁺ concentration was calculated based on the fluorescence ratio and Ca²⁺ calibration curve. To measure intracellular ROS production, cells were loaded with 10 μ M CM-DCFH diacetate for 15 min at room temperature in the dark. The coverslip was affixed to a culture chamber and perfused with the same culture medium without phenol red. Under each experimental condition about 15 single myocytes were imaged with an Attofluor imaging system (Atto Instruments), and CM-DCF fluorescence was measured at excitation wavelength of 480 nm and emission wavelength of 520 nm.

Protein synthesis

Protein synthesis in cultured myocytes was assayed by [³H]phenylalanine incorporation as we described before [4,9]. Cells were cultured in 12-well plates, treated with glucose oxidase and other agents as indicated for 44 h, and continued for an additional 4 h in the presence of 0.5 μ Ci of [³H]phenylalanine/well before total protein and [³H]phenylalanine incorporation were assayed.

Northern blot

Total RNA was isolated using TRI reagent according to the manufacturer's protocol. Northern blot was done as previously described [9]. Routinely, about 20 μ g of total RNA was subjected to gel electrophoresis, transferred to a Nytran membrane, UV-immobilized, and hybridized to ³²P-labeled probes. Autoradiograms obtained at -70°C were scanned with a Bio-Rad densitometer. The relative amount of RNA in each sample was normalized by using GAPDH mRNA as an internal control.

Assay for ⁸⁶Rb⁺ uptake by cardiac myocytes

The initial rate of Rb⁺ uptake through the Na⁺/K⁺-ATPase was measured in the presence or absence of 1 mM ouabain as we described before [9]. Monensin (20 μ M) was added to the medium for 10 min at 37°C prior to the initiation of the ⁸⁶Rb⁺ uptake assay to assure that the maximal capacity of active uptake was measured.

Measurement of ERK1/2 activity

Activation of ERK1/2 in cultured myocytes was determined as we previously reported [29] by Western blot using a monoclonal anti-phospho-ERK1/2 antibody that detects ERK1/2 only when they are phosphorylated at Thr202 and Tyr204. The same blots were stripped and probed with a polyclonal antibody recognizing the total amount of ERK1/2 to account for equal loading. The blots were scanned with a Bio-

Rad densitometer to quantify phospho- and total ERK1/2 signals.

Labeling of cell surface Na/K-ATPase by biotinylation

Cell surface protein biotinylation was performed as described before [30]. Proteins bound to the ImmunoPure immobilized streptavidin-agarose beads were eluted and then resolved on SDS-PAGE followed by immunoblotting.

Confocal microscopy

Cell growth, immunostaining, and confocal microscopy were performed as described before with little modification [6]. Briefly, after treatment, myocytes were fixed with 2% paraformaldehyde for 10 min at room temperature and then permeabilized in 0.3% Triton X-100 for 10 min. Cells were blocked by Image-iT FX signal enhancer (Molecular Probes) for 30 min. For cell size and actin staining, cells were incubated with Alexa 488-conjugated Phalloidin (Molecular Probes) for 2 h at room temperature. For Na/K-ATPase endocytosis, cells were incubated with anti-Na/K-ATPase- α_1 monoclonal antibody (Upstate Biotechnology, 1:50) and followed by incubation with Alexa 488-conjugated secondary antibodies. The coverslips were mounted with ProLong Gold antifade reagent (Molecular Probes). Confocal image was captured by a Leica TCS SP2 spectral confocal scanner and a Leica DMIRE2 microscope (Leica, Mannheim, Germany) equipped with a 63 \times oil immersion objective. Visualization and analysis were performed using Leica confocal microscope system software.

Statistics

Data are given as the mean \pm SE. Statistical analysis was performed using the Student's *t* test, and significance was accepted at $p < 0.05$. Each presented immunoblot is representative of the similar results of at least three separate experiments.

Results

Addition of glucose oxidase generates chronic ROS stress in cardiac myocytes

Since increases in ROS play a pivotal role in ouabain-induced hypertrophic growth, it is of interest to extend our studies on ouabain and determine if chronic exposure of cardiac myocytes to elevated ROS regulates cardiac growth and gene expression in a pattern similar to that of ouabain. In order to achieve this goal, we added glucose oxidase to the culture medium to continuously generate H_2O_2 . It is known that glucose oxidase produces H_2O_2 in the presence of glucose [31]. As expected, addition of this enzyme to the culture medium caused a sustained rise in intracellular ROS (Fig. 1A). Addition of catalase (50 U/ml) completely blocked glucose oxidase-induced increases in intracellular ROS, indicating that increases in intracellular ROS were due to glucose oxidase-catalyzed




Fig. 1. Stimuli-induced changes in intracellular ROS. (A) Cardiac myocytes were treated with indicated concentrations of glucose oxidase (G.O.), H_2O_2 , ouabain (Oua), or catalase (50 U/ml) plus glucose oxidase (Cat.+G.O.) for various times. (B) Cardiac myocytes were treated with glucose oxidase and/or ouabain for various times. The stimuli-treated cells were then loaded with 10 μ M CM-DCFH diacetate and assayed for the fluorescence intensities of the oxidized probe as described under Experimental procedures. The value of every time point was calibrated by the value of the control cells. Each value is the mean \pm SE of determinations of 15 single cells in four independent experiments.

H_2O_2 production. The effects of glucose oxidase on intracellular ROS levels were dose and time dependent. Glucose oxidase-induced increases in ROS reached maximal levels after 4 h and lasted for at least 12 h. In comparison, exposure of myocytes to a bolus 10–25 μ M H_2O_2 caused a much faster increase in intracellular ROS. However, intracellular ROS decreased significantly from the peak after 1 h H_2O_2 exposure and approached the control level after 2–4 h of incubation (Fig. 1A). When compared to added bolus H_2O_2 , 100 μ M ouabain increased intracellular ROS to levels similar to that induced by 2 μ M H_2O_2 , while 3 mU/ml glucose oxidase raised intracellular ROS to that produced by 5 μ M H_2O_2 (data not shown). Clearly, like ouabain, the addition of glucose oxidase (but not a bolus H_2O_2) to culture medium could cause a sustained increase in intracellular ROS levels. Moreover, the treatment with ouabain and glucose oxidase simultaneously caused an additive effect on ROS generation (Fig. 1B). Therefore, we used this glucose oxidase model in the following studies to investigate whether exposure of cultured cardiac myocytes to chronic ROS stress induced hypertrophic growth.

Chronic ROS stress stimulates hypertrophic growth in cardiac myocytes

To address whether an increase in ROS alone is sufficient to cause hypertrophic growth in cardiac myocytes, we first measured the effects of glucose oxidase on the expression of hypertrophic marker genes using Northern blot analysis. A characteristic feature of hypertrophic growth in adult cardiac muscle is the activation of a so-called “fetal program” that consists of the recapitulation of a pattern of cardiac-specific gene expression of the prenatal heart. For example, the rodent skACT is expressed in the fetal ventricle, and then replaced by the adult isoform, cardiac α -actin, during postnatal development. Many hypertrophic stimuli including ouabain increase the expression of skACT as well as ANF in cultured myocytes [32]. In addition, these stimuli also reduce the expression of the α_3 subunit of the Na^+/K^+ -ATPase [9]. As depicted in Figs. 2A and 2B, glucose oxidase, like ouabain, caused a dose-dependent increase in the expression of both skACT and ANF mRNA. In addition, it also reduced the expression of the Na^+/K^+ -ATPase α_3 mRNA. The glucose oxidase-induced decrease in α_3 mRNA and increases in skACT and ANF mRNA were completely




Fig. 2. Effects of glucose oxidase on the expression of hypertrophic marker genes. Myocytes were treated with different concentrations of glucose oxidase (G.O.) alone or with 10 mM NAC or 50 U/ml catalase for 12 h, and total RNA was isolated and assayed for cardiac gene expression using Northern blot. Panel A show a representative Northern blot. Lane 1, control; Lane 2, 1.5 mU/ml glucose oxidase; Lane 3, 3.0 mU/ml glucose oxidase; Lane 4, 3.0 mU/ml glucose oxidase plus NAC. Panel B shows quantitative data. Values were normalized to those of corresponding GAPDH measured on the same blot and expressed relative to control value of one. Each value is the mean \pm SE of three experiments.

blocked by either catalase or NAC. To further test whether increases in ROS alone are sufficient to induce hypertrophic growth, cells were exposed to different concentrations of glucose oxidase and assayed for morphological changes and [^3H]phenylalanine incorporation. As shown in Fig. 3, glucose oxidase clearly increased the cell size and stimulated actin reorganization in the cultured myocytes as other hypertrophic stimuli [26,33]. Consistently, glucose oxidase also stimulated protein synthesis in a dose-dependent manner (Fig. 4). It is important to note that under the same experimental conditions, a bolus H_2O_2 at 10 μM or lower showed no effect on protein synthesis (data not shown), while a bolus 25 μM H_2O_2 caused a significant decrease in total protein content. When cell viability was determined, 10 μM H_2O_2 had no effect, while 25 μM H_2O_2 caused a statistically significant loss (about 15%) of cell viability after 24 h incubation. On the other hand, ouabain up to 100 μM and glucose oxidase up to 3 mU/ml showed no effect on cell viability after 48 h of exposure. These findings, taken together with those in Fig. 1, clearly showed potential pitfalls associated with use of a high concentration of bolus H_2O_2 in studying long-term effects of ROS signaling. A low dose of H_2O_2 , as shown in Fig. 1A, would be destroyed quickly by the cell antioxidant system so that it could not exhibit a significant effect on cell growth whereas a high dose might cause cell toxicity. In short, the above data indicate that increases in intracellular ROS alone were sufficient to stimulate hypertrophic growth in cultured neonatal rat cardiac myocytes.

ROS-induced cardiac hypertrophy shared signal pathways similar to those activated by ouabain binding to the Na^+/K^+ -ATPase

The Na^+/K^+ -ATPase serves as an important signal transducer. Binding of ouabain to the Na/K -ATPase activates multiple pathways including increases in ROS [9], intracellular Ca^{2+} [3], and activation of ERK1/2 [7,34]. There is evidence that increases in intracellular Ca^{2+} and subsequent activation of the calcineurin/NF-AT pathway as well as the stimulation of ERK1/2 play an important role in ouabain and other stimuli-induced hypertrophic growth [4,35]. Because glucose oxidase and ouabain caused a similar pattern of changes in gene expression and cell growth, the following experiments were performed to explore whether glucose oxidase affected intracellular Ca^{2+} and ERK1/2 activity in cardiac myocytes. Activation of ERK1/2 was measured by Western blot using an anti-phospho-ERK antibody. As depicted in Fig. 5, addition of 3 mU/ml glucose oxidase increased phosphorylation levels of ERK1 (upper 44-kDa band) and 2 (42-kDa band) in a time-dependent manner. This time course of ERK activation correlated with the rise in intracellular ROS (Fig. 1A). When intracellular Ca^{2+} was measured in fura-2-loaded myocytes, glucose oxidase caused a gradual rise in intracellular Ca^{2+} as shown in Fig. 6. The maximal effects were observed after the cells were exposed to glucose oxidase for 2–4 h and the intracellular Ca^{2+} remained elevated for at least 12 h.

To investigate the functionality of the ERK1/2 pathway in glucose oxidase-induced hypertrophic growth, myocytes were




Fig. 3. Effects of glucose oxidase on sarcomere reorganization and cell size. Myocytes were treated with glucose oxidase or/and ouabain for 24 h, fixed, and then stained with Alexa 488-conjugated phalloidin as described under Experimental procedures. Scale bar = 10 µm. A representative experiment is shown and the same experiments were repeated three times.

pretreated with a MEK inhibitor (PD98059) for 30 min and then exposed to glucose oxidase. As shown in Fig. 7, PD98059 completely abolished the effect of glucose oxidase on protein synthesis. To test the role of a rise in intracellular Ca^{2+} and the resulting activation of the calcineurin/NF-AT pathway [36–38], myocytes were preloaded with 10 µM BAPTA-AM and then exposed to glucose oxidase. Although BAPTA significantly decreased basal protein synthesis, it did not block glucose oxidase-induced protein synthesis (Fig. 8). Because ouabain signals via the activation of Src [39] and increases in ROS are known to stimulate Src [40,41], we also tested the role of Src family kinase. As depicted in Fig. 7, inhibition of Src by PP2 completely blocked glucose oxidase-induced increases in protein synthesis.

Cross talk between Na^+/K^+ -ATPase and ROS

We have shown that partial inhibition of Na^+/K^+ -ATPase by ouabain increased ROS production in cardiac myocytes [9]. Since our previous work showed that ROS had the potential to interact with the cardiac Na^+/K^+ -ATPase [35], we reasoned that the Na^+/K^+ -ATPase could serve as a target for ROS to transmit the signals to downstream effectors, thus contributing to ROS-induced changes in gene expression and cardiac hypertrophy in




Fig. 4. Effects of glucose oxidase and ouabain on protein synthesis. Myocytes were treated with glucose oxidase (G.O.) or/and ouabain (Oua) for 48 h and then assayed for [³H]phenylalanine incorporation as indicated under Experimental procedures. Values are means ± SE of four experiments. **p* < 0.05, ***p* < 0.01 compared to control.

cultured neonatal cardiac myocytes. To address this issue, we first determined whether ouabain and ROS have additive effects on cell signaling, growth, and gene expression. Consistent with the findings illustrated in Fig. 1B, ouabain potentiated the effects of glucose oxidase on gene expression (data not shown) and protein synthesis (Fig. 4), supporting the notion that glucose and ouabain share similar pathways of cardiac hypertrophy.

To further test whether ROS cross talk with the ouabain-activated pathways, we measured the effects of ROS on the cardiac Na^+/K^+ -ATPase activity. It showed that sustained increases in ROS stress produced by glucose oxidase could cause a time-dependent inhibition of Na^+/K^+ -ATPase (Fig. 9). Inhibition of the Na^+/K^+ -ATPase by glucose oxidase correlated well with the increases in intracellular ROS concentration. The maximal inhibition (about 40%) occurred after 4 h of glucose oxidase exposure. Since the enzyme is significantly inhibited by




Fig. 5. Effect of glucose oxidase on activation of ERK1/2. Myocytes were exposed to glucose oxidase (3 mU/ml) for the indicated times and assayed for ERK1/2 activation as described under Experimental procedures. (A) Representative immunoblots. Membranes were detected with anti-phosphorylated ERK1/2 antibody and then stripped and reprobed by an anti-ERK1/2 pan antibody that recognizes total ERK1/2. (B) Quantitative data from three independent experiments. The phosphorylated ERK1/2 values were normalized against the total ERK1/2 signal. This ratio was then expressed relative to control value of one. Values are means ± SE **p* < 0.05 vs control.




Fig. 6. Long-term effects of glucose oxidase and ouabain on intracellular Ca^{2+} . Myocytes were treated with either glucose oxidase and/or ouabain for various times and then loaded with fura-2 AM. Myocyte fura-2 signals were measured as described under Experimental procedures. Each value is the mean \pm SE determinations on 40 cells in four independent experiments.

glucose oxidase at the concentration that stimulates hypertrophic growth, it is reasonable to suggest that Na^+/K^+ -ATPase is involved in ROS signaling in cardiac myocytes. Previously, we showed that the activation of signal transduction by ouabain stimulated endocytosis of Na^+/K^+ -ATPase [30,42]. To test whether ROS can activate the Na^+/K^+ -ATPase-mediated pathways, and result in the endocytosis, we measured the surface α_1 subunit of Na^+/K^+ -ATPase by cell surface biotinylation assay after myocytes were exposed to glucose oxidase for various times. As shown in Fig. 10, in response to 3 mU/ml glucose oxidase (6 h), the surface Na^+/K^+ -ATPase α_1 subunit decreased by about 40%. To further confirm these findings, we immunostained the cultured myocytes with Na^+/K^+ -ATPase α_1 subunit antibody after the cells were treated with glucose oxidase for 2 h. Confocal microscopy images showed that increases in ROS stress were able to induce internalization of the plasmalemmal Na^+/K^+ -ATPase α_1 subunit in cardiac myocytes (Fig. 11). As expected, this induction of internaliza-




Fig. 7. Effects of PP2 and PD98059 on glucose oxidase-induced protein synthesis. Myocytes were pretreated with 0.5 μ M PP2 or 5 μ M PD98059 for 30 min and then exposed to glucose oxidase (3 mU/ml) for 48 h and assayed for $[^3H]$ phenylalanine incorporation as described under Experimental procedures. Values are means \pm SE of four experiments. * p < 0.05 compared to control, # p < 0.05 compared to glucose oxidase.




Fig. 8. Effects of BAPTA on glucose oxidase-induced protein synthesis. Myocytes were loaded with 10 μ M BAPTA-AM for 30 min and then exposed to glucose oxidase for 48 h. $[^3H]$ Phenylalanine incorporation was performed as in Fig. 4. Values are means \pm SE of four experiments. * p < 0.05 compared to control, # p < 0.05 compared to glucose oxidase.

tion was completely blocked when catalase (50 U/ml) was added to the culture medium prior to the addition of glucose oxidase (data not shown).

Discussion

In this report we demonstrated that the addition of glucose oxidase to glucose-containing medium continuously produced H_2O_2 , which resulted in a sustained increase in intracellular ROS concentration. The sustained, but not transient, increases in intracellular ROS alone were sufficient to activate hypertrophic growth pathways, leading to phenotypic changes similar to those induced by ouabain and other hypertrophic stimuli in cultured neonatal rat cardiac myocytes. Significantly, ROS may cross talk with the pathways emanated from the signaling Na^+/K^+ -ATPase.

ROS are known to play an important role in the pathogenesis of cardiovascular diseases [10–13,43]. Recent studies have




Fig. 9. Time-dependent inhibition of ouabain-sensitive $^{86}Rb^+$ uptake by H_2O_2 and glucose oxidase. Myocytes cultured in 12-well plates were treated with 25 μ M H_2O_2 or 3 mU/ml glucose oxidase for various times. Ouabain-sensitive $^{86}Rb^+$ uptakes were performed as described under Experimental procedures. Each value is the mean \pm SE of four experiments.




Fig. 10. Glucose oxidase reduces surface expression of Na⁺/K⁺-ATPase α₁ subunit. Myocytes were exposed to glucose oxidase (3 mU/ml) for the indicated times and assayed for surface expression by biotinylation as described under Experimental procedures. (A) A representative Western blot analysis of biotinylated α₁. (B) Quantitative data are presented as mean ± SE of four individual experiments. * *p* < 0.05 vs control.

demonstrated that ROS is an important second messenger that regulate various cellular functions including cell growth [10,23,44]. We and others have shown that activation of many receptors including Na⁺/K⁺-ATPase elevates intracellular ROS [22,26,45–47]. Blocking the rise in intracellular ROS reduces the agonist (e.g., ouabain)-induced activation of signaling pathways as well as hypertrophic growth in cultured cardiac myocytes [9]. However, it was difficult to study the direct effect of ROS on various signaling pathways and on cell growth because the addition of a bolus H₂O₂ does not mimic chronic hypertrophic stimuli-induced ROS stress. This is well-illustrated in Figs. 1 and 4. While the low doses of bolus H₂O₂ did not affect cell growth because they were rapidly removed by the cellular antioxidant system, high doses of ROS were toxic to myocytes. It was reported that the addition of glucose oxidase to culture medium could continuously produce H₂O₂ for over 24 h in the presence of glucose [31]. Taking advantage of these findings we assessed whether glucose oxidase could produce a sustained and modest ROS stress as ouabain did in cultured myocytes. Indeed, glucose oxidase raised intracellular ROS levels in a time- and dose-dependent manner in the cultured myocytes. By varying the enzyme concentration we were able to titrate intracellular ROS to concentrations comparable to those induced by ouabain. Thus, we have developed a protocol that allows us to study the effect of changes in intracellular ROS on cellular signaling and function.

Interestingly, when the effects of ROS on gene expression and cell growth were assessed, we found that a sustained increase in ROS alone was sufficient to stimulate protein synthesis and actin reorganization in cultured cardiac myocytes (Figs. 3 and 4). Interestingly, inhibition of Src and ERK1/2, but not Ca²⁺, pathways abrogated the effects of glucose oxidase on protein synthesis (Figs. 7 and 8). Like many other hypertrophic stimuli [9,20,21], glucose oxidase-induced cell growth was accompanied by changes in expression of several hypertrophic

marker genes (Fig. 2). Thus, chronic ROS stress alone can function as hypertrophic stimuli that may act through the protein kinases or cellular receptors such as Na⁺/K⁺-ATPase in induction of hypertrophic growth in cardiac myocytes. In principle, there are three types of cardiac hypertrophy: normal growth, physiologic hypertrophy, and pathologic hypertrophy [48]. Typically, stimuli of G protein-coupled receptors induced pathologic hypertrophy, while exercise and growth factors activated physiologic hypertrophy. Although it is inconclusive whether ROS-induced hypertrophic growth is physiological or pathological based on the data presented in this work, it is likely that ROS stress contributes to the development of maladaptive hypertrophy [10,48].

In principle, ROS can regulate cell growth by directly modulating the functions of the signaling proteins such as tyrosine phosphatases and kinases in cardiac myocytes. Alternatively, ROS can affect the receptors that are important for generating hypertrophic growth signals. The Na⁺/K⁺-ATPase is an important receptor for both endogenous and exogenous cardiotoxic steroids such as ouabain [49,50]. Binding of ouabain to the Na⁺/K⁺-ATPase stimulates protein tyrosine phosphorylation, resulting in subsequent activation of multiple protein kinase cascades that are known to play important roles in the control of cell growth [4,6,8,39,51,52]. It also increases intracellular Ca²⁺ and ROS production, resulting in hypertrophic growth in cardiac myocytes [3,9]. Interestingly, ROS are known to directly modify the structure of the Na⁺/K⁺-ATPase [53,54], and the effects of various ROS on Na⁺/K⁺-ATPase activity as well as ouabain binding sites are well documented in cardiac myocytes [55–57]. Thus, it is conceivable that at least some of the glucose oxidase effects on cell growth could be mediated through the Na⁺/K⁺-ATPase. Indeed, we showed here that increase in ROS stress acted as ouabain and stimulated endocytosis of the Na⁺/K⁺-ATPase (Figs. 10 and 11), which resulted in a significant inhibition of Na⁺/K⁺-ATPase activity [42]. This could be important because the Na⁺/K⁺-ATPase may serve as a molecular target for ROS.




Fig. 11. Glucose oxidase induces Na⁺/K⁺-ATPase endocytosis. Myocytes were treated with 3 mU/ml glucose oxidase for 2 h and then fixed and probed for α₁ subunit of Na⁺/K⁺-ATPase as described under Experimental procedures. Images were scanned by Leica confocal microscope. A representative experiment is shown. Bar: 10 μm. A majority of the α₁ subunit was detected in the plasma membrane in control myocytes whereas a significant portion of the α₁ subunit was moved to intracellular compartments after myocytes were treated with glucose oxidase.

Signaling through the Na⁺/K⁺-ATPase by ROS may generate second messengers such as intracellular Ca²⁺. Indeed, as depicted in Fig. 6, ROS-induced changes in intracellular Ca²⁺ appeared to be correlated to its effect on the Na⁺/K⁺-ATPase. In addition, ouabain and ROS exhibited the additive effect on intracellular Ca²⁺, suggesting a common target for both stimuli. This notion was further supported by the findings showing that ouabain potentiated the effects of ROS on protein synthesis (Fig. 4) and gene expression (data not shown). Thus, ROS may not only regulate protein kinase cascade directly but also affect membrane receptors such as Na⁺/K⁺-ATPase, thus reinforcing its effect on hypertrophic signaling pathways. To this end, it is worth noting that ouabain binding to the Na⁺/K⁺-ATPase stimulates ROS production in a Src and Ras-dependent manner in cardiac myocytes [3,9]. Therefore, it is likely that ouabain-induced ROS could further affect the receptor, creating a signal amplification loop. Clearly, this hypothesis remains to be tested experimentally.

Acknowledgments

We thank Jennifer Kalisz for her assistance in the preparation of neonatal cardiac myocytes. This work was supported by National Institutes of Health Grants HL-36573 and HL-63238 awarded by the National Heart, Lung, and Blood Institute, United States Public Health Service, Department of Health and Human Services.

References

- [1] Skou, J. C.; Esmann, M. The Na,K-ATPase. *J. Bioenerg. Biomembr.* **24**: 249–261; 1992.
- [2] Lingrel, J. B.; Kuntzweiler, T. Na⁺,K⁺-ATPase. *J. Biol. Chem.* **269**: 19659–19662; 1994.
- [3] Liu, J.; Tian, J.; Haas, M.; et al. Ouabain interaction with cardiac Na⁺/K⁺-ATPase initiates signal cascades independent of changes in intracellular Na⁺ and Ca²⁺ concentrations. *J. Biol. Chem.* **275**:27838–27844; 2000.
- [4] Huang, L.; Li, H.; Xie, Z. Ouabain-induced hypertrophy in cultured cardiac myocytes is accompanied by changes in expression of several late response genes. *J. Mol. Cell. Cardiol.* **29**:429–437; 1997.
- [5] Peng, M.; Huang, L.; Xie, Z.; et al. Partial inhibition of Na⁺/K⁺-ATPase by ouabain induces the Ca²⁺-dependent expressions of early-response genes in cardiac myocytes. *J. Biol. Chem.* **271**:10372–10378; 1996.
- [6] Liu, L.; Mohammadi, K.; Aynafshar, B.; et al. Role of caveolae in signal-transducing function of cardiac Na⁺/K⁺-ATPase. *Am. J. Physiol. Cell Physiol.* **284**:C1550–C1560; 2003.
- [7] Mohammadi, K.; Kometiani, P.; Xie, Z.; et al. Role of protein kinase C in the signal pathways that link Na⁺/K⁺-ATPase to ERK1/2. *J. Biol. Chem.* **276**:42050–42056; 2001.
- [8] Haas, M.; Askari, A.; Xie, Z. Involvement of Src and epidermal growth factor receptor in the signal-transducing function of Na⁺/K⁺-ATPase. *J. Biol. Chem.* **275**:27832–27837; 2000.
- [9] Xie, Z.; Kometiani, P.; Liu, J.; et al. Intracellular reactive oxygen species mediate the linkage of Na⁺/K⁺-ATPase to hypertrophy and its marker genes in cardiac myocytes. *J. Biol. Chem.* **274**:19323–19328; 1999.
- [10] Dhalla, A. K.; Hill, M. F.; Singal, P. K. Role of oxidative stress in transition of hypertrophy to heart failure. *J. Am. Coll. Cardiol.* **28**:506–514; 1996.
- [11] Giordano, F. J. Oxygen, oxidative stress, hypoxia, and heart failure. *J. Clin. Invest.* **115**:500–508; 2005.
- [12] Chien, K. R. Stress pathways and heart failure. *Cell* **98**:555–558; 1999.
- [13] Griendling, K. K.; Sorescu, D.; Lassegue, B.; et al. Modulation of protein kinase activity and gene expression by reactive oxygen species and their role in vascular physiology and pathophysiology. *Arterioscler. Thromb. Vasc. Biol.* **20**:2175–2183; 2000.
- [14] Takemoto, M.; Node, K.; Nakagami, H.; et al. Statins as antioxidant therapy for preventing cardiac myocyte hypertrophy. *J. Clin. Invest.* **108**:1429–1437; 2001.
- [15] Griendling, K. K.; Ushio-Fukai, M. Reactive oxygen species as mediators of angiotensin II signaling. *Regul. Pept.* **91**:21–27; 2000.
- [16] Sorescu, D.; Griendling, K. K. Reactive oxygen species, mitochondria, and NAD(P)H oxidases in the development and progression of heart failure. *Congest. Heart Fail.* **8**:132–140; 2002.
- [17] Takano, H.; Zou, Y.; Hasegawa, H.; et al. Oxidative stress-induced signal transduction pathways in cardiac myocytes: involvement of ROS in heart diseases. *Antioxid. Redox Signal* **5**:789–794; 2003.
- [18] Sawyer, D. B.; Siwik, D. A.; Xiao, L.; et al. Role of oxidative stress in myocardial hypertrophy and failure. *J. Cell. Cardiol.* **34**:379–388; 2002.
- [19] Yamamoto, M.; Yang, G.; Hong, C.; et al. Inhibition of endogenous thioredoxin in the heart increases oxidative stress and cardiac hypertrophy. *J. Clin. Invest.* **112**:1395–1406; 2003.
- [20] Priyadarshi, S.; Valentine, B.; Han, C.; et al. Effect of green tea extract on cardiac hypertrophy following 5/6 nephrectomy in the rat. *Kidney Int.* **63**:1785–1790; 2003.
- [21] Kennedy, D. J.; Vetteth, S.; Periyasamy, S. M.; et al. Central role for the cardiotoxic steroid marinobufagenin in the pathogenesis of experimental uremic cardiomyopathy. *Hypertension* **47**:488–495; 2006.
- [22] Mollnau, H.; Wendt, M.; Szocs, K.; et al. Effects of angiotensin II infusion on the expression and function of NAD(P)H oxidase and components of nitric oxide/cGMP signaling. *Circ. Res.* **90**:E58–E65; 2002.
- [23] Amin, J. K.; Xiao, L.; Pimental, D. R.; et al. Reactive oxygen species mediate alpha-adrenergic receptor-stimulated hypertrophy in adult rat ventricular myocytes. *J. Mol. Cell. Cardiol.* **33**:131–139; 2001.
- [24] Chen, Q. M.; Tu, V. C.; Purdon, S.; et al. Molecular mechanisms of cardiac hypertrophy induced by toxicants. *Cardiovasc. Toxicol.* **1**:267–283; 2001.
- [25] Thannickal, V. J.; Fanburg, B. L. Reactive oxygen species in cell signaling. *Am. J. Physiol. Lung Cell Mol. Physiol.* **279**:L1005–L1028; 2000.
- [26] Xu, F. P.; Chen, M. S.; Wang, Y. Z.; et al. Leptin induces hypertrophy via endothelin-1-reactive oxygen species pathway in cultured neonatal rat cardiomyocytes. *Circulation* **110**:1269–1275; 2004.
- [27] Pimentel, D. R.; Amin, J. K.; Xiao, L.; et al. Reactive oxygen species mediate amplitude-dependent hypertrophic and apoptotic responses to mechanical stretch in cardiac myocytes. *Circ. Res.* **89**:453–460; 2001.
- [28] Tu, V. C.; Bahl, J. J.; Chen, Q. M. Distinct roles of p42/p44(ERK) and p38 MAPK in oxidant-induced AP-1 activation and cardiomyocyte hypertrophy. *Cardiovasc. Toxicol.* **3**:119–133; 2003.
- [29] Kometiani, P.; Li, J.; Gnudi, L.; et al. Multiple signal transduction pathways link Na⁺/K⁺-ATPase to growth-related genes in cardiac myocytes. The roles of Ras and mitogen-activated protein kinases. *J. Biol. Chem.* **273**:15249–15256; 1998.
- [30] Liu, J.; Liang, M.; Liu, L.; et al. Ouabain-induced endocytosis of the plasmalemmal Na⁺/K⁺-ATPase in LLC-PK1 cells requires caveolin-1. *Kidney Int.* **67**:1844–1854; 2005.
- [31] Lai, C. C.; Peng, M.; Huang, L.; et al. Chronic exposure of neonatal cardiac myocytes to hydrogen peroxide enhances the expression of catalase. *J. Mol. Cell. Cardiol.* **28**:1157–1163; 1996.
- [32] Karns, L. R.; Kariya, K.; Simpson, P. C. M-CAT, CARG, and Sp1 elements are required for alpha 1-adrenergic induction of the skeletal alpha-actin promoter during cardiac myocyte hypertrophy. Transcriptional enhancer factor-1 and protein kinase C as conserved transducers of the fetal program in cardiac growth. *J. Biol. Chem.* **270**:410–417; 1995.
- [33] Morisco, C.; Zebrowski, D. C.; Vatner, D. E.; et al. Beta-adrenergic cardiac hypertrophy is mediated primarily by the beta(1)-subtype in the rat heart. *J. Mol. Cell. Cardiol.* **33**:561–573; 2001.
- [34] Mohammadi, K.; Liu, L.; Tian, J.; et al. Positive inotropic effect of ouabain on isolated heart is accompanied by activation of signal pathways that link Na⁺/K⁺-ATPase to ERK1/2. *J. Cardiovasc. Pharmacol.* **41**:609–614; 2003.
- [35] Xie, Z. J.; Wang, Y. H.; Askari, A.; et al. Studies on the specificity of the effects of oxygen metabolites on cardiac sodium pump. *J. Mol. Cell. Cardiol.* **22**:911–920; 1990.

- [36] Hiroi, Y.; Hiroi, J.; Kudoh, S.; et al. Two distinct mechanisms of angiotensin II-induced negative regulation of the mitogen-activated protein kinases in cultured cardiac myocytes. *Hypertens. Res.* **24**: 385–394; 2001.
- [37] Vega, R. B.; Bassel-Duby, R.; Olson, E. N. Control of cardiac growth and function by calcineurin signaling. *J. Biol. Chem.* **278**:36981–36984; 2003.
- [38] Guo, J.; Zhan, S.; Somers, J.; et al. Decrease in Density of INa is in the common final pathway to heart block in murine hearts overexpressing calcineurin. *Am. J. Physiol. Heart Circ. Physiol.*; 2006.
- [39] Tian, J.; Cai, T.; Yuan, Z.; et al. Binding of Src to Na⁺/K⁺-ATPase forms a functional signaling complex. *Mol. Biol. Cell* **17**:317–326; 2006.
- [40] Sato, H.; Sato, M.; Kanai, H.; et al. Mitochondrial reactive oxygen species and c-Src play a critical role in hypoxic response in vascular smooth muscle cells. *Cardiovasc. Res* **67**:714–722; 2005.
- [41] Ushio-Fukai, M.; Zuo, L.; Ikeda, S.; et al. cAbl tyrosine kinase mediates reactive oxygen species- and caveolin-dependent AT1 receptor signaling in vascular smooth muscle: role in vascular hypertrophy. *Circ. Res.* **97**:829–836; 2005.
- [42] Liu, J.; Kesiry, R.; Periyasamy, S. M.; et al. Ouabain induces endocytosis of plasmalemmal Na⁺/K⁺-ATPase in LLC-PK1 cells by a clathrin-dependent mechanism. *Kidney Int.* **66**:227–241; 2004.
- [43] Dhalla, N. S.; Temsah, R. M.; Netticadan, T. Role of oxidative stress in cardiovascular diseases. *J. Hypertens.* **18**:655–673; 2000.
- [44] Chen, Q. M.; Tu, V. C.; Wu, Y.; et al. Hydrogen peroxide dose dependent induction of cell death or hypertrophy in cardiomyocytes. *Arch. Biochem. Biophys.* **373**:242–248; 2000.
- [45] Hool, L. C. Reactive oxygen species in cardiac signalling: from mitochondria to plasma membrane ion channels. *Clin. Exp. Pharmacol. Physiol.* **33**:146–151; 2006.
- [46] Itoh, S.; Ding, B.; Bains, C. P.; et al. Role of p90 ribosomal S6 kinase (p90RSK) in reactive oxygen species and protein kinase C beta (PKC-beta)-mediated cardiac troponin I phosphorylation. *J. Biol. Chem.* **280**:24135–24142; 2005.
- [47] Thannickal, V. J.; Hassoun, P. M.; White, A. C.; et al. Enhanced rate of H₂O₂ release from bovine pulmonary artery endothelial cells induced by TGF-beta 1. *Am. J. Physiol.* **265**:L622–L626; 1993.
- [48] Dorn, G. W., II; Force, T. Protein kinase cascades in the regulation of cardiac hypertrophy. *J. Clin. Invest.* **115**:527–537; 2005.
- [49] Scheiner-Bobis, G.; Schoner, W. A fresh facet for ouabain action. *Nat. Med.* **7**:1288–1289; 2001.
- [50] Schoner, W. Endogenous cardiac glycosides, a new class of steroid hormones. *Eur. J. Biochem.* **269**:2440–2448; 2002.
- [51] Liu, J.; Shapiro, J. I. Endocytosis and signal transduction: basic science update. *Biol. Res. Nurs.* **5**:117–128; 2003.
- [52] Liu, L.; Abramowitz, J.; Askari, A.; et al. Role of caveolae in ouabain-induced proliferation of cultured vascular smooth muscle cells of the synthetic phenotype. *Am. J. Physiol. Heart Circ. Physiol.* **287**:H2173–H2182; 2004.
- [53] Huang, W. H.; Wang, Y.; Askari, A.; et al. Different sensitivities of the Na⁺/K⁺-ATPase isoforms to oxidants. *Biochim Biophys. Acta* **1190**: 108–114; 1994.
- [54] Zolotarjova, N.; Ho, C.; Mellgren, R. L.; et al. Different sensitivities of native and oxidized forms of Na⁺/K⁺-ATPase to intracellular proteinases. *Biochim. Biophys. Acta* **1192**:125–131; 1994.
- [55] Jamme, I.; Petit, E.; Divoux, D.; et al. Modulation of mouse cerebral Na⁺, K⁺-ATPase activity by oxygen free radicals. *Neuroreport* **7**:333–337; 1995.
- [56] Kako, K.; Kato, M.; Matsuoka, T.; et al. Depression of membrane-bound Na⁺-K⁺-ATPase activity induced by free radicals and by ischemia of kidney. *Am. J. Physiol.* **254**:C330–C337; 1988.
- [57] Kim, M. S.; Akera, T. O₂ free radicals: cause of ischemia-reperfusion injury to cardiac Na⁺-K⁺-ATPase. *Am. J. Physiol.* **252**:H252–H257; 1987.