

Review

Emerging perspectives in store-operated Ca^{2+} entry: Roles of Orai, Stim and TRP

Jeremy T. Smyth, Wayne I. DeHaven, Bertina F. Jones, Jason C. Mercer, Mohamed Trebak,
Guillermo Vazquez, James W. Putney Jr.*

*Laboratory of Signal Transduction, National Institute of Environmental Health Sciences, National Institutes of Health,
Department of Health and Human Services, Research Triangle Park, NC 27709, USA*

Received 5 July 2006; accepted 30 August 2006
Available online 5 September 2006

Abstract

Depletion of intracellular Ca^{2+} stores induces Ca^{2+} influx across the plasma membrane through store-operated channels (SOCs). This store-operated Ca^{2+} influx is important for the replenishment of the Ca^{2+} stores, and is also involved in many signaling processes by virtue of the ability of intracellular Ca^{2+} to act as a second messenger. For many years, the molecular identities of particular SOC, as well as the signaling mechanisms by which these channels are activated, have been elusive. Recently, however, the mammalian proteins STIM1 and Orai1 were shown to be necessary for the activation of store-operated Ca^{2+} entry in a variety of mammalian cells. Here we present molecular, pharmacological, and electrophysiological properties of SOC, with particular focus on the roles that STIM1 and Orai1 may play in the signaling processes that regulate various pathways of store-operated entry.

Published by Elsevier B.V.

Keywords: Calcium channel; TRP channel; Stim1; Orai1; Calcium-release-activated-calcium current; Store-operated channel; Capacitative calcium entry; Calcium oscillation

Ca^{2+} is an essential second messenger in virtually all cell types, and it regulates a wide spectrum of events from neurotransmission to gene transcription. Cytoplasmic Ca^{2+} can be elevated in cells either by release from internal stores located in the endoplasmic reticulum (ER) or by entry across the plasma membrane [9]. In many cases, intracellular Ca^{2+} release and Ca^{2+} entry occur in concert with one another, and thus both

routes of increasing the intracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) are often necessary for proper signaling to occur.

A major mechanism of Ca^{2+} release involves the phosphoinositide (PI) pathway, whereby plasma membrane receptors of the G-protein-linked or tyrosine kinase families activate specific phospholipase C (PLC) isoforms, resulting in the cleavage of phosphatidylinositol 4,5-bisphosphate (PIP_2) into diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP_3). DAG is most recognizable as an activator of protein kinase C (PKC), but can also play a more direct role in regulating Ca^{2+} signal generation. IP_3 is a critical Ca^{2+} releasing agent in the cell, and it achieves this by activating the IP_3 receptor (IP_3R) located in the ER membrane; binding of IP_3 to its receptor results in the receptor/channel opening and release of Ca^{2+} from the ER into the cytoplasm [7].

Activation of the PI pathway also results in Ca^{2+} entry by 2 important mechanisms. First, second messengers generated downstream of activation of the PI pathway, such as DAG, arachidonic acid (AA), and Ca^{2+} itself [11], can directly activate Ca^{2+} entry channels in the plasma membrane; these channels

Abbreviations: $[\text{Ca}^{2+}]_i$, intracellular Ca^{2+} concentration; 2-APB, 2-aminoethoxydiphenylborane; AA, arachidonic acid; ARC, arachidonate-regulated channel; CCE, capacitative Ca^{2+} entry; CIF, Ca^{2+} influx factor; DAG, diacylglycerol; ER, endoplasmic reticulum; EYFP, enhanced yellow fluorescent protein; I_{CRAC} , Ca^{2+} release-activated Ca^{2+} current; IP_3 , inositol 1,4,5-trisphosphate; IP_3R , inositol 1,4,5-trisphosphate receptor; OAG, 1-oleoyl-2-acetyl-sn-glycerol; PI, phosphoinositide; PIP_2 , phosphatidylinositol 4,5-bisphosphate; PKC, protein kinase C; PLA_2 , phospholipase A_2 ; PLC, phospholipase C; ROC, receptor-operated channel; SAM, sterile-alpha motif; SCID, severe combined immunodeficiency; SERCA, sarco/endoplasmic reticulum Ca^{2+} -ATPase; SOC, store-operated channel; TRP, transient receptor potential

* Corresponding author.

E-mail address: putney@niehs.nih.gov (J.W. Putney).

are referred to as receptor-operated channels (ROCs). Second, the release of Ca^{2+} from the ER induced by IP_3 leaves the ER in a state of Ca^{2+} depletion; this depletion of ER Ca^{2+} activates plasma membrane Ca^{2+} entry [98], a process originally termed capacitative calcium entry (CCE) [128], through channels generally referred to as store-operated channels (SOCs). This review will focus on Ca^{2+} entry through SOCs, covering the most recent findings in the activation mechanisms of SOCs, the pharmacological, electrophysiological, and molecular properties of SOCs, and physiological and pathophysiological roles for SOCs. For more information on other modes of receptor-regulated entry, the reader is referred to recent reviews [11,140].

1. Overview of store-operated entry

By definition, SOCs are activated exclusively as a result of depletion of intracellular Ca^{2+} stores [86]. Experimentally, entry through SOCs can be distinguished from other modes of Ca^{2+} entry by employing protocols that achieve ER Ca^{2+} depletion independently of activation of other signaling pathways. Paramount to the idea that SOC activation relies exclusively on store depletion is the fact that the ER must be able to communicate Ca^{2+} store depletion to SOCs within the plasma membrane in some way.

The existence of SOCs was first postulated in 1986 [98]. Since then, the molecular nature of the process, i.e., the channels responsible for the Ca^{2+} influx and the signaling mechanisms that result in activation of the channels, has remained enigmatic. It is important to note that store-operated entry does not necessarily refer to a single mechanism of Ca^{2+} entry, nor does SOC necessarily refer to a single Ca^{2+} entry channel; instead, any channel that can be shown to exhibit Ca^{2+} store-dependent activity can be referred to as a SOC. The most studied and best characterized SOC current is the Ca^{2+} release-activated Ca^{2+} current, or I_{CRAC} , which was first characterized in mast cells [45,46], but has been measured in a variety of cells ranging from T and B-lymphocytes to hepatocytes [85]. I_{CRAC} is a non-voltage-gated current, and is notable for its very high selectivity for Ca^{2+} , as indicated by its strong inward rectification, highly positive reversal potential, and low conductance [85,86,97]; other store-operated entry currents that have been characterized generally exhibit less stringent ionic selectivities and a variety of conductances [85,86]. The mechanism by which I_{CRAC} is activated, including the molecular identity of the CRAC channel, has long been the most sought-after signaling paradigm in the store-operated entry field. For some time now, a significant effort has been made to determine whether or not members of the transient receptor potential (TRP) family of ion channels are molecular components of the CRAC channel. Particular focus has concentrated on the canonical TRPs (TRPCs), since it has been shown that these ion channels are activated downstream of PLC activation and, in some cases, can operate as SOCs (see Section 5.1); however, as of yet, no TRPC channel has passed the ultimate test—that is, the ability to fully recapitulate the electrophysiological and pharmacological properties of I_{CRAC} . Nevertheless, it is important to note that TRP channels may function in certain physiological contexts as SOCs, despite the

fact that this activity is distinct from I_{CRAC} . Now, the quest to uncover the molecular identity of the I_{CRAC} channel and its mechanism of activation has come to an important new turning point with the recent discoveries that two proteins, STIM1 and Orai1, play obligatory roles in the activation of I_{CRAC} . As will be discussed in Section 4, accumulating evidence indicates that STIM1 acts as a sensor of ER store content, while Orai1 may be the I_{CRAC} channel itself.

Regardless of its form, whether it be via I_{CRAC} or another SOC, store-operated Ca^{2+} influx accomplishes several critical functions within the cell. First, Ca^{2+} that enters the cell via the SOC pathway replenishes the ER Ca^{2+} stores following a release event, thus maintaining the ability of the ER to release Ca^{2+} into the cytoplasm in response to subsequent Ca^{2+} releasing stimuli. This is most notable in cells that respond to activation of the PI pathway with $[\text{Ca}^{2+}]_i$ oscillations, since ablation of Ca^{2+} entry by removing extracellular Ca^{2+} or by other pharmacological or molecular means will often times preclude a cell's ability to maintain $[\text{Ca}^{2+}]_i$ oscillations (see Section 6). Second, Ca^{2+} concentrations within the ER must be maintained at sufficient levels in order for the organelle to carry out many of its fundamental functions. Thus, chronic depletion of ER Ca^{2+} , as would occur in the absence of store-operated Ca^{2+} entry, can influence ER-dependent processes such as protein folding and trafficking, the ER stress response, and apoptosis [20]. And third, it should be noted that Ca^{2+} that enters the cell via the store-operated pathway first accesses the cytoplasm before entering the ER and, in many cases, results in sustained elevation in $[\text{Ca}^{2+}]_i$ levels. This is significant in that for some signaling events, in particular T-lymphocyte activation, it is the increase in cytoplasmic Ca^{2+} that results from store-operated entry, as opposed to that which results from Ca^{2+} released from the ER, that is responsible for signaling [54]. Thus, store-operated entry can influence many aspects of cell biology, and it is vital that we understand the molecular compositions of all forms of store-operated entry so that we may fully appreciate its contribution to normal physiological as well as pathophysiological states of human health.

2. Pharmacology of store-operated entry

The use of pharmacological inhibitors of sarco/endoplasmic reticulum Ca^{2+} -ATPase (SERCA) pumps, which are responsible for pumping Ca^{2+} into the lumen of the ER from the cytoplasm, has been invaluable in the study of SOC function and regulation. These compounds, which include thapsigargin, cyclopiazonic acid and di-*tert*-butylhydroquinone, achieve Ca^{2+} store depletion by allowing a constitutive passive leak of Ca^{2+} from the ER to occur unchecked due to the inability of the inhibited SERCA pumps to replenish the ER [21]. Further, ER depletion can be effected by dialysis of cells with high concentrations of Ca^{2+} buffers such as EGTA or BAPTA, which effectively reduce the free Ca^{2+} concentration in the cytosol available to the SERCA pumps. Finally, direct buffering of ER Ca^{2+} with membrane permeant *N,N,N',N'*-tetrakis(2-pyridylmethyl)ethylene diamine (TPEN) reduces the free Ca^{2+} concentration in the ER and activates store-operated entry. Hence, the pharmacological tools

with which SOC activation can be revealed are quite extensive; however, these tools do not allow for manipulation of the entry process itself since they act on mechanisms that are upstream of channel activation. In order to define roles for store-operated entry in physiological processes, it is important to have available tools by which one can predictably and specifically antagonize the entry process. Unfortunately, the SOC field has chronically been plagued by a lack of pharmacological inhibitors that specifically target store-operated Ca^{2+} influx.

Currently available pharmacological inhibitors of the store-operated entry pathway lack optimal specificity. Indeed, many agents such as econazole and SK&F-96365 that are known to inhibit SOCs were also noted to inhibit other ion channels, particularly voltage-dependent channels, within the same concentration ranges [34,99]. Another drug called 2-aminoethoxydiphenylborane (2-APB) described originally as an IP_3R antagonist [69], was shown later to be a direct inhibitor of SOCs, having no effect on voltage-dependent Ca^{2+} entry channels or arachidonic acid-activated Ca^{2+} channels [2,15,96]. However, 2-APB was later shown to interfere with a wide variety of transport mechanisms including SERCA pumps, potassium channels and mitochondrial Ca^{2+} efflux (for review see [86]). In addition, 2-APB has been found to activate TRPV1, TRPV2 and TRPV3, members of the heat-gated vanilloid TRP (TRPV) family of cation channels [23,47]. Despite these various effects of 2-APB unrelated to store-operated entry, the drug remains an important tool in the study of Ca^{2+} influx. Store-operated entry and I_{CRAC} in particular have the unique characteristics of being potentiated by low concentrations of 2-APB (1–5 μM), whereas they are inhibited by concentrations greater than 10 μM [66,96]. And although the mechanism by which 2-APB exerts these effects on SOCs is unknown, this biphasic sensitivity to 2-APB is often used as a “fingerprint” that helps define the store-operated nature of a particular Ca^{2+} influx process.

There are other reagents that block store-operated entry but also block voltage-dependent Ca^{2+} entry channels with equal efficiencies. These drugs include L561582 [29,99], SC38249 [25,99], LU52396 [24,99] and tetrandine [53,99]. Other drugs such as cytochrome P-450 inhibitors or reagents that inhibit enzymes of arachidonic acid metabolism (cyclooxygenase and lipoxygenase inhibitors) are also known to inhibit store-operated entry (for review see [99]). However, these drugs are not specific and their exact action on store-operated entry is not clear.

The most reliable inhibitors of SOCs remain trivalents such as La^{3+} and Gd^{3+} . The latter when used at concentrations in the low micromolar (1–5 μM) range has proven to be a specific blocker of store-operated entry in many cell types without affecting arachidonic acid-activated Ca^{2+} entry or the Ca^{2+} entry pathway mediated by receptor-activated TRPC channels [17,63,135]. However, this inorganic inhibitor precipitates with divalent anions and thus cannot be used *in vivo*. Its use is therefore restricted to *in vitro* studies using cultured cells in media with controlled compositions.

The quest for specific inhibitors of the various SOCs expressed in human cells is clearly in its infancy and new drugs that interfere with SOC activation are likely to be promising

tools for treatment of human diseases. The recent discovery of STIM1 and Orai1 as important determinants in store-operated entry is an important breakthrough toward our understanding of the exact molecular identity and mechanism of activation of SOCs. With further appreciation of the molecular makeup and mechanism of activation of store-operated Ca^{2+} influx, we can expect future developments of reagents that specifically target this pathway.

3. Activation mechanisms of store-operated entry

At present there are 2 general hypotheses for the activation mechanism of various store-operated entry pathways and of I_{CRAC} in particular: (i) action of a diffusible messenger that is released from the ER to activate plasma membrane channels upon ER Ca^{2+} depletion, and (ii) coupling between proteins within the ER membrane with Ca^{2+} influx channels in the plasma membrane [85,86]. In addition, there is speculation that the activation mechanism may involve insertion of channels into the plasma membrane [153] although some mechanism of initiating the signaling from the ER would still be necessary. There is evidence in support of these ideas in the literature; however, much of this evidence is indirect and discrepancies abound [86]. Direct proof of any of these activation mechanisms as they relate to store-operated entry will likely only come as a result of the identification of the molecular nature of individual SOC channels.

3.1. Diffusible messenger

The diffusible messenger mechanism was recently the topic of an extensive review [14] and will thus only be briefly addressed here. The theory of a diffusible factor responsible for activating store-operated entry was first implied by Putney and colleagues [127], and evidence for the existence of such a factor was first demonstrated by Randriamampita and Tsien [103]. In this latter work the authors were successful in isolating an acid-extractable factor from activated Jurkat T-lymphocytes, referred to as Ca^{2+} influx factor (CIF), that was capable of activating Ca^{2+} influx when applied to several different cell types. Importantly, further biochemical purification of CIF led to the resolution of a factor that was only effective at evoking Ca^{2+} influx when microinjected into cells [50,131], and thus acted upon an intracellular target. Another significant boost to the CIF story came when it was shown that a biochemically and functionally similar factor could also be isolated from *Saccharomyces cerevisiae* in which the Ca^{2+} -ATPase gene *pmr1* is mutated, resulting in depletion of intracellular Ca^{2+} stores [26]; thus, CIF production is apparently not specific to a single cell type or species. This study was also the first to show a positive effect of CIF on the development of I_{CRAC} [26]. The mechanism by which CIF activates Ca^{2+} influx remains unclear, although recent work suggests a role for Ca^{2+} -independent phospholipase A_2 (iPLA₂). It was first shown by pharmacological and molecular approaches that inhibition of iPLA₂ activity results in suppression of SOC activity [41,112,120]. Smani et al. [119]

then demonstrated that CIF is able to activate store-operated entry by displacing an inhibitory calmodulin from iPLA₂, and that lysophospholipid generated by iPLA₂ is responsible for directly activating SOC channels. However, the exact mechanism by which CIF activates SOC and whether it constitutes the physiological activation signal for I_{CRAC} awaits the identification of the molecular entity of CIF.

3.2. Conformational coupling

The conformational coupling hypothesis, first proposed by Irvine [48] and further developed by Berridge [8], draws analogy to excitation–contraction coupling in skeletal muscle whereby the dihydropyridine-sensitive Ca²⁺ channel in the plasma membrane directly couples with the type 1 ryanodine receptor in the sarcoplasmic reticulum. In the case of store-operated entry, it was proposed that the IP₃R in the ER membrane directly links with putative Ca²⁺ entry channels in the plasma membrane, and that a conformational change of the IP₃R upon ER Ca²⁺ depletion signals the opening of SOC channels. This hypothesis was revised somewhat by Patterson et al. [88] into what has come to be known as the secretion-like coupling model. This secretion-like coupling model differs from the original conformational coupling model in that the former allows for separation between IP₃Rs and SOCs when stores are full, and in response to store depletion the ER, or specific subdomains of the ER, move close to the plasma membrane and allow the interaction to occur between the IP₃R and SOCs. The most direct evidence for such a mechanism comes from studies demonstrating that stabilization of a cortical ring of actin with agents such as jasplakinolide or calyculin A prevents the activation of store-operated entry [88,111], the idea being that this ring of actin prevents the movement of the ER toward the plasma membrane. However, a subsequent exhaustive study in RBL-1 cells failed to demonstrate any effects of cytoskeletal alterations on the development of I_{CRAC} [2]; a similar lack of dependence of store-operated entry on the cytoskeleton had also been previously demonstrated in NIH-3T3 cells [106]. Thus, cell-type differences appear to be important considerations in the conclusions drawn from experiments that evaluate the effects of the cytoskeleton on store-operated entry. Evidence in favor of conformational coupling between the IP₃R and SOC channels was also gleaned from early studies showing that 2-APB, thought at the time to be a specific IP₃R inhibitor [69], prevents thapsigargin-induced Ca²⁺ entry in HEK293 cells [64]. It was subsequently shown that this drug also inhibits the development of I_{CRAC} in Jurkat and RBL cells [2,15,16,96]. However, these effects of 2-APB on store-operated entry can no longer be taken as proof of involvement of the IP₃R, since it was later shown that 2-APB has direct inhibitory effects on CRAC channels independently of antagonism of the IP₃R [2,15,96]. Most significantly, it was shown that 2-APB is equally effective at inhibiting store-operated entry and I_{CRAC} in a DT40 B-cell line in which all 3 isoforms of the IP₃R had been genetically ablated, as compared to its inhibitory effect in wild type DT40 cells [15,65]. And in fact the finding that Ca²⁺

entry and I_{CRAC} develop normally in these IP₃R knockout cells [125] provides the strongest evidence yet to indicate that interaction of Ca²⁺ influx channels with the IP₃R is not required for the activation of I_{CRAC}. However, this does not preclude the idea that other proteins within the ER membrane may conformationally couple with CRAC channels and achieve I_{CRAC} activation. Significantly, it has recently been shown that STIM1 may act as such a factor, since it has been shown that STIM1 is able to sense Ca²⁺ store levels within the ER lumen, and undergoes a major relocalization toward the plasma membrane in response to Ca²⁺ store depletion [59,110,123,157], as will be addressed in Section 4. Thus, STIM1 may act as the conformational coupler between the ER and CRAC channels, an idea that has been addressed in a recent review [43], and the conformational coupling literature may now need to be readdressed in the context of STIM1 (discussed below).

3.3. Vesicle fusion

The vesicle fusion hypothesis postulates that vesicles containing functional SOCs fuse with the plasma membrane upon store depletion, resulting in plasma membrane insertion of active channels. Evidence in support of this mechanism comes from experimental strategies whereby I_{CRAC} or Ca²⁺ influx are measured under conditions in which exocytosis or vesicle fusion should be impaired. For example, it was shown that pharmacological or molecular interference with SNAP-25, a member of the soluble N-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) complex, antagonizes store-operated currents in *Xenopus* oocytes [153]. It was also shown in HEK 293 cells that botulinum and tetanus neurotoxins, which disrupt SNAP-25 and vesicle-associated membrane protein 2 (VAMP-2), respectively, significantly reduced store-depletion-activated Ca²⁺ entry [1]. However, another study challenged the interpretation of these findings [114]. An important caveat of all studies examining the roles of vesicle trafficking is that it is difficult to distinguish from such studies a role for induced exocytosis in response to store depletion, as is required by the fusion hypothesis, from disruption of constitutive exocytosis which is necessary to maintain the normal complement of membrane components for SOC function. Furthermore, it will be impossible to directly demonstrate store-driven exocytosis of a CRAC channel until the molecular identity of this channel is known. Interestingly, in the case of TRPC channels, several reports have demonstrated that agonists that activate channel activity also result in exocytotic insertion of these channels [10,22,81,84,118], thus further supporting the idea that exocytosis may play an important role in Ca²⁺ entry mechanisms. However, others have failed to observe such increases in membrane expression of TRPC channels in response to activating maneuvers [115,121]. And, as will be discussed in Section 5, it remains unclear whether TRP channels mediate physiological store-operated entry, and in all of the instances cited above, TRP channels were activated by means other than store depletion.

4. STIM1 and Orai1: Molecular components of I_{CRAC}

Through the use of RNAi technology, two proteins have recently been identified that are necessary for store-operated entry and I_{CRAC} in particular. The first of these proteins was identified simultaneously by two separate groups using RNAi in *Drosophila* S2 cells [110] and in human HeLa cells [59]. Each group found that when the messages for the proteins Stim in *Drosophila* or STIM1 in mammalian cells were targeted using siRNA, cells were unable to activate Ca^{2+} entry following store depletion using thapsigargin.

STIM1 was identified in limited screens using bioinformatics to identify genes with probable ion channel or signaling functions. However, the second protein, termed Orai or CRACM, bears no resemblance or homology to any other functional protein; thus it is not surprising that its discovery came somewhat later, resulting from RNAi screens of the entire *Drosophila* genome [33,143,156]. In the first publication describing this protein, its identification was facilitated by the finding that this particular gene was also the genetic cause of a rare form of severe combined immunodeficiency (SCID) [32,33]. The identification of these two proteins has revitalized the quest to understand the mechanisms controlling I_{CRAC} .

4.1. STIM1

Roos et al. performed a large scale RNAi screen in *Drosophila* S2 cells by targeting any protein with a transmembrane domain, including the TRP channel family, as well as any protein that had been previously described as playing a role in store-operated entry [110]. Of 170 genes targeted, only Stim was found to be needed for store-operated entry and activation of I_{CRAC} in S2 cells. In a parallel study Liou et al. targeted a total of 2304 proteins. They identified two mammalian Stim homologues, STIM1 and STIM2, whose knockdown reduced store-operated entry [59]. It should be noted that when Roos et al. applied siRNAs against STIM1 and STIM2 in Jurkat cells, STIM1 was essential for store-operated entry whereas loss of STIM2 had no effect [110]. Perhaps the discrepancy between these two results is due to off target effects of the siRNA used in the Liou study, which was prepared by incubating the mRNA for the respective proteins with the protein Dicer and then treating cells with the processed RNA [59]. It is conceivable that one or more pieces of the dicer-processed STIM2 RNA bore significant homology to the STIM1 message. These reports also showed that while treatment of cells with siRNA against STIM1 abolishes store-operated entry, it has no apparent effect on the intracellular Ca^{2+} store content or the resting cytosolic Ca^{2+} concentration [59,110,123].

STIM1 is a type 1a single-pass transmembrane protein [68,146]. It contains several conserved domains, including a sterile-alpha motif (SAM), a coiled coil region, and an EF-hand domain. Interestingly, neither the SAM nor the EF-hand domains bear full resemblance to their respective consensus domains. The SAM domain contains two N-linked glycosylation sites, making it the first glycosylated SAM domain to be identified [68,146]. EF-hand domains normally occur in pairs;

however STIM1 only has one. There is some evidence to suggest that STIM1 can homodimerize or heterodimerize with STIM2, and this could bring together the EF-hands from separate molecules to form an intermolecularly paired EF-hand domain [146]. Due to the presence of the putative EF-hand domain and the unlikely prospect of a channel protein containing only one transmembrane domain, STIM1 is thought to be the Ca^{2+} store-sensing component of the I_{CRAC} pathway as opposed to the CRAC channel itself. In agreement with this prediction, mutations in acidic residues within the Ca^{2+} binding pocket of the EF-hand domain of STIM1, which presumably lower the affinity for Ca^{2+} , produce constitutive Ca^{2+} entry that is independent of store depletion [59,157]. Additionally, wild-type STIM1 expressed as a fusion with enhanced yellow fluorescent protein (EYFP-STIM1) co-localized at least partially with a marker for ER but rapidly rearranged into punctate structures following store depletion [59]. EF-hand mutants of EYFP-STIM1 that confer constitutive Ca^{2+} entry also appear constitutively in punctate structures, indicating that Ca^{2+} binding is important for the localization of STIM1 [59]. However, without physical evidence of actual Ca^{2+} binding, it must be considered that the EF-hand domain of STIM1 might not bind Ca^{2+} and could serve another purpose.

There is some debate as to the cellular localization of STIM1 following store depletion. Several groups have reported detection of STIM1 on the cell surface [68,123,146,157]. However, others have failed to detect any STIM1 in the plasma membrane [59,73]. Additionally, while Zhang et al. [157] reported a significant increase in plasma membrane STIM1 following store depletion, Spassova et al. [123] reported no such increase. Both Manji et al. [68] and Williams et al. [146] reported plasma membrane STIM1, but store depletion was not evaluated in these studies because STIM1 was not known to be involved in store-operated entry at the time. Thus, a portion of cellular STIM1 may be inserted into the plasma membrane, but it is unclear if those STIM1 molecules are important for activation of I_{CRAC} .

Although it is clear that STIM1 is a necessary component of I_{CRAC} and store-operated entry, the mechanism by which it acts is still a mystery. Three preliminary models have been proposed to explain how STIM1 might work based upon its cellular localization patterns. All three models have in common that STIM1 is located in the ER membrane prior to store depletion. The first model suggests that STIM1 in the ER complexes with STIM1 in the plasma membrane following store depletion [110]. This model of course requires that some STIM1 is found in the plasma membrane, and that this pool of STIM1 is necessary for activation of I_{CRAC} . The second model contends that most or all STIM1 is confined to the ER membrane prior to store depletion. After store depletion however, STIM1 (which is presumably no longer bound to Ca^{2+}) is somehow transported to and inserted into the plasma membrane [157]. It is not clear what mechanism would be involved in the transport of STIM1 to the plasma membrane, as accumulating evidence suggests that the most likely mechanism, vesicle trafficking, is not involved in I_{CRAC} [2,86]. Both the first and second models are supported by

cell surface biotinylation experiments that detected STIM1 [123,157]. However, these data must be interpreted with caution since STIM1 may have been co-precipitated with a cell surface protein such as the channel itself rather than being directly biotinylated. In studies employing transfection of HEK293 cells with STIM1 and Orai1 (discussed in the following section), no STIM1 was detected in the plasma membrane by antibody binding, despite the generation of huge STIM1- and Orai1-dependent I_{CRAC} -like currents [73]. These latter findings favor a third model which places STIM1 only in the ER membrane [59]. Following store depletion, the punctae seen with EYFP-STIM1 would be very close to the cell surface, but not actually inserted into the plasma membrane. In this scenario the cytosolic C-terminus of STIM1 would interact with additional components of the I_{CRAC} pathway, such as the channels themselves, when brought into the punctae. This model does not exclude the possibility that STIM1 under certain circumstances can be found in the plasma membrane. It does indicate however that this putative plasma membrane pool of STIM1 is not essential for activation of I_{CRAC} and store-operated entry.

4.2. Orai1

The second protein component of I_{CRAC} that has been identified was named Orai after the Greek mythological characters the Orai which are the keepers of the gates of heaven [33]. In a report following close thereafter, the name CRACM (for CRAC-Membrane) was suggested [143]. We will use the Orai designation in deference to the first published description of this protein. In addition to Orai1, there are two other Orai family members, Orai2 and Orai3, expressed in mammals [33]. Flies and worms express only one Orai homolog, and this protein most closely resembles the mammalian Orai1 [33]. Orai1 was first identified from genetic linkage analysis performed to identify mutations contributing to a rare form of severe SCID characterized by a lack of I_{CRAC} activity, combined with a genome-wide screen in *Drosophila* to search for genes necessary for nuclear translocation of the Ca^{2+} -dependent transcription factor NFAT [32,33]. Orai was identified as essential for Ca^{2+} entry in *Drosophila*, and the role of human Orai1 was confirmed by sequencing the Orai1 gene from the SCID patients as well as 21 of their relatives. In addition to the two SCID patients, 13 of the relatives had lower I_{CRAC} activity in their T cells and were predicted to be heterozygous carriers of the mutation [33]. Importantly, all 13 heterozygous family members as well as the 2 SCID patients carried the same arginine to tryptophan mutation at amino acid position 91 in Orai1 [33]. This mutation was confirmed to block I_{CRAC} by its failure to rescue the lack of Ca^{2+} entry and I_{CRAC} activity when transfected into T cells and fibroblasts from the SCID patients, whereas transfection of the wild-type version did rescue [33].

Orai1, as well as the related Orai2 and Orai3, are predicted to be membrane spanning proteins with 4 transmembrane domains [33,143,156]. Unlike STIM1, Orai1 apparently resides only at the plasma membrane [33,143]. With the exception of the transmembrane segments, these proteins have no other readily identifiable modular domains. Shortly following the report of

the role of Orai1 in the human SCID patients, Vig et al. also identified *Drosophila* Orai as an essential component of I_{CRAC} , although they called it CRACM1 [143]. In addition to Orai, Vig et al. [143] identified another membrane component that they termed CRACM2; however there is no human ortholog of CRACM2 and so it appears not to be a conserved component of the I_{CRAC} pathway. Subsequently, several reports have demonstrated a remarkable amplification of I_{CRAC} (50–100 fold) when Orai1 is coexpressed with STIM1 [90,122,156]. This amplification is not seen when STIM1 alone is transfected into cells. When Orai1 alone is transfected, there is a clear inhibition of I_{CRAC} in Jurkat [90], RBL [122] cells and HEK293 cells [73] (but see [90]). These findings illustrate several important points. First, they demonstrate that STIM1 and Orai1 are the only components necessary to produce large I_{CRAC} currents when overexpressed, indicating that there are no other rate-limiting components of the system (unless unknown endogenous components are upregulated by the overexpression conditions). Interestingly, the I_{CRAC} currents observed when STIM1 and Orai1 are coexpressed do not completely recapitulate all of the canonical characteristics of I_{CRAC} . Most notably, the amplified currents have low permeability for Ba^{2+} and Sr^{2+} [143], both of which are known to pass readily through native I_{CRAC} channels although to a lesser extent than Ca^{2+} [163]. This could indicate that other protein components are found in native I_{CRAC} channels, such as the other Orai family members, that are not necessary for activation and Ca^{2+} permeation, but that modulate some of the other pore properties. Second, the finding that Orai1 overexpression alone results in inhibition of I_{CRAC} suggests Orai1 interacts with at least two other similarly limiting components of the SOC pathway, perhaps STIM1 itself. Thus when excess Orai1 is present, the statistical likelihood of combining in an appropriate stoichiometric fashion with these other limiting components is reduced, reducing the number of correctly assembled signaling complexes [101]. Third, the observation that overexpression of these two proteins produces such a dramatic increase in I_{CRAC} current density argues (but does not prove) that Orai1 may be the I_{CRAC} pore-forming unit. If Orai1 were merely another part of the activation machinery then overexpression of STIM1 plus Orai1 might result in a small increase in current density, but only to the point at which the endogenous channels are all in the open state. Especially in HEK293 cells which have nearly undetectable endogenous I_{CRAC} , the likelihood of the cells expressing enough endogenous channels to accommodate such large increases in current density seems unlikely.

Mercer et al. [73] reported that Orai2 exhibits properties similar to Orai1; i.e., overexpression of Orai2 alone resulted in inhibition of store-operated Ca^{2+} entry in HEK293 cells, whereas its coexpression with STIM1 resulted in significantly augmented entry and I_{CRAC} . The magnitude of the currents with Orai2 were somewhat less than for Orai1, however. Orai3, on the other hand, did not synergistically enhance Ca^{2+} entry when coexpressed with STIM1. However, Orai3 was shown to restore Ca^{2+} entry in cells in which entry was reduced due to RNAi-mediated knockdown of endogenous Orai1, demonstrating that Orai3 is able to function in the Ca^{2+} permeation pathway. Thus,

in transient transfection experiments, the rank order of efficacy of Orai family members appeared to be Orai1>Orai2>Orai3 [73].

Our developing appreciation of the critical roles that STIM1 and Orai1 play in the regulation of I_{CRAC} will undoubtedly revolutionize the field of store-operated entry. However, it must be kept in mind that I_{CRAC} may be only one species of store-operated entry that operates physiologically in mammalian cells. Therefore, it is important that we continue to examine other store-operated entry pathways, such as those potentially mediated by TRP channels and other, possibly undiscovered channels. Additionally, it will be interesting to determine whether STIM and Orai isoforms influence store-operated entry pathways that are distinct from I_{CRAC} .

5. TRP proteins as SOCs

For over a decade, members of the superfamily of TRP proteins, particularly those from the mammalian TRPC family, have been considered molecular candidates of SOCs. Based on amino acid homology, the TRP superfamily includes at present seven families: the TRPC (Canonical; TRPC1–TRPC7), the TRPV (Vanilloid; TRPV1–6), and the TRPM (Melastatin; TRPM1–8) families, and other more distantly related members, like the TRPML (Mucolipin) and TRPP (Polycystin) families, which although not yet well characterized, have received increasing attention due to their association with several inherited human diseases. More recently, the TRPA (Ankyrin) group, with only one mammalian member, and the TRPN (no mechanoreceptor potential C) group, only found in *Caenorhabditis*, *Drosophila* and zebra fish, have been incorporated into the TRP superfamily (reviewed in [89,102]). Recognition of the product of the *Drosophila* *trp* gene as the Ca^{2+} permeable channel underlying light-induced, PLC-dependent Ca^{2+} entry in fly photoreceptor cells (reviewed in [77]), was rapidly followed by cloning of several mammalian *trp* counterparts, TRPC1 and TRPC3 being the first to be cloned and expressed in cell lines [13]. Immediately thereafter, realization that ectopically expressed TRPC proteins also make Ca^{2+} permeable cation channels, in most cases activated downstream of PLC stimulation, brought about the belief that these proteins could function as the long-sought SOCs. Consequently, most, if not all studies aimed to elucidate the molecular nature of SOCs focused on mammalian TRPC proteins. However, rather than illuminating, the resultant literature provided controversial evidence for each of the TRPC proteins as well as for TRPV6 and TRPM3, either in favor of or refuting a role as SOCs. To illustrate this controversy, an overview of the major findings regarding the role of TRPs as SOCs is provided below (for a thorough discussion, see [86,102,136,140]).

5.1. TRPCs

The TRPC family contains the mammalian genes most closely related to the *Drosophila* *trp* gene [145,159]. Based on structural similarities the TRPC family is divided into four subgroups: TRPC1; TRPC2; TRPC3, 6 and 7; and TRPC4 and

5. TRPC3, 6 and 7 not only share a high degree of amino acid identity but also functional and regulatory similarities. A structural, and possibly functional relationship also exists between TRPC4 and 5. TRPC2, a pseudo gene in humans and old world primates, forms functional channels in mouse, rat and other mammalian species [161,162]. The reader is referred to a recent review for a comprehensive discussion of structure–function relationships of TRPC channel proteins [140].

The notion that TRPC1 and 4 may form, or be part of SOCs is supported mainly by two different lines of experimental evidence. First, ectopic expression of TRPC1 or 4 in several cell lines increases endogenous store-operated entry induced by either PLC-dependent or pharmacologically-induced depletion of ER Ca^{2+} stores [36,61,91,144,160]. Second, the use of antisense constructs directed against TRPC1 or 4, as well as genetic disruption of the corresponding genes, has efficiently reduced store-operated entry in a variety of cell types [18,19,61,126,132,150,152]. In some instances, those protocols produced a significant reduction of the archetypical store-operated current I_{CRAC} [35,78,93], leading to the conclusion that TRPC1 or 4 are part of native CRAC channels. However, the question remains if the decrease in I_{CRAC} was directly caused by an actual reduction in TRPC1 or 4 protein levels, rather than being the result of compensatory changes in the expression of CRAC channel subunits or regulatory proteins. It is noteworthy that the focused and whole genome screens described above that brought the roles of Stim1 and Orai1 to light [33,59,110,143,156] apparently failed to detect any TRP channels. Moreover, neither TRPC1 nor TRPC4 recapitulate the biophysical properties expected for CRAC channels when ectopically expressed in cell lines. Despite the evidence mentioned above, the role of TRPC4 in store-operated entry has not yet received as much consensus as for TRPC1 (see [3,95] and references therein). This is mainly due to the controversy raised by several studies showing that ectopic expression of mouse TRPC4 in diverse cell lines results in the appearance of PLC-regulated cation channels whose activity is completely independent of Ca^{2+} store depletion [80,113,158]. Also, in HEK293 cells, a human epithelial cell line, ectopically expressed human TRPC4 rendered constitutively active channels not regulated by agonist or Ca^{2+} store-depletion [71]. Evidence for store-operated regulation of mouse TRPC5 is limited, and published results are controversial (cf. [92] and [82]). In most instances TRPC5 activation, although downstream of receptor stimulation of PLC, seems not to be related to Ca^{2+} store-depletion [95,113]. A more recent report however, showed that at the single cell level TRPC5 may possibly be activated by a variety of signals, including depletion of Ca^{2+} stores [155]. It is conceivable that the disparate channel behavior, in terms of sensitivity to store depletion, could be influenced by the expression conditions (see below).

Ectopic expression of a splice variant of mouse TRPC2, designated TRPC2A, was shown to enhance store-operated Ca^{2+} entry in monkey kidney fibroblast-like cells [138]. However, a different lab cloned two additional splice variants (named TRPC2 α and TRPC2 β) [44], and showed that neither these variants nor TRPC2A were able to form channels operated by

Ca²⁺ store-depletion when expressed in HEK293 cells, in conflict with a role for mouse TRPC2 as a SOC. Conversely, native TRPC2 has been reported to take part in endogenous store-operated entry in Chinese hamster ovary cells [37] and spermatid cells [49].

TRPC3 constitutes an inveterate example of the existing controversy as to whether or not a particular TRPC channel can function in a store-operated manner. Some of the confusion regarding the ability of TRPC3 to function in a store-operated mode stems from the fact that, when ectopically expressed, TRPC3 constitutes a channel endowed with significant constitutive activity. This makes it difficult to distinguish a genuine store-dependent effect from mere enhanced constitutive activity, particularly in experiments where Ca²⁺ entry is monitored by means of fluorescent Ca²⁺ indicators (discussed in [100,136]). Despite compelling evidence supporting the view that the signaling to TRPC3 is downstream of receptor-dependent activation of PLC but in a manner not related to Ca²⁺ store depletion (reviewed in [136,140]), studies by Kiselyov et al. [51,52] advocating that TRPC3 channels are indeed store-operated, challenged that view. These studies showed that stable expression of TRPC3 in HEK293 cells results in a non-selective cation channel activated either by stimulation of a Gq-coupled muscarinic receptor or by pharmacological depletion of Ca²⁺ stores, in both instances in an IP₃R-dependent fashion. In contrast, Trebak et al. [134] showed that in two different clones of TRPC3-expressing HEK293 cells, including the one used in the studies by Kiselyov et al. [51,52], TRPC3 behaves as a ROC independently of Ca²⁺ store depletion or the IP₃R. A possible explanation for these disparate results regarding the store-dependence/independence of TRPC3 channels derives from recent studies by Vazquez et al. [139,142], showing that in the avian B-lymphocyte DT40 cell line ectopic expression of TRPC3 resulted in a channel whose ability to function in either receptor- or store-operated modes was clearly related to channel expression level (discussed in detail in [101]). Store-operation was observed only at very low channel expression levels, whereas conditions that resulted in higher levels of TRPC3 protein expression gave rise to the receptor-regulated mode of TRPC3 activation.

TRPC7 was originally shown to function as a PLC-regulated channel when expressed in HEK293 cells [83], presumably through PLC-derived DAG, but subsequent evidence indicated that it can also function as a SOC [107]. Notably, recent work revealed that these apparently conflicting findings stem from the ability of TRPC7 to operate in either store-operated or receptor-activated modes depending, as for TRPC3, upon the protein expression level [56]. In those studies, Lievremont et al. [56] showed that transient expression of TRPC7 in HEK293 cells rendered a DAG-activated channel, but when expressed stably, it behaved as a SOC. In an attempt to determine which of these two coupling modes corresponds to the physiological function of endogenously expressed TRPC7, a follow up study [57] utilized targeted homologous recombination to knock out TRPC7 in DT40 B-lymphocytes, known to express endogenously both store-operated and non-store-operated pathways [87,125,141]. In what represented the first demonstration of a physiological

function for native TRPC7, it was found that knockout of TRPC7 in DT40 B-lymphocytes did not significantly affect I_{CRAC} [57]. Rather, in the avian B cell line, TRPC7 functions as a PLC-regulated, DAG-activated non-selective cation channel. It is possible, however, that native TRPC7 may function as a SOC in other cell types, for instance, when SOC currents exhibit properties reminiscent of TRPCs (see for example [137]).

As of this writing, available evidence indicates that ectopic expression of TRPC6, either of mouse or human origin, results in a non-selective cation channel whose activation is linked to PLC-derived products but is independent of Ca²⁺ store depletion (reviewed in [136,140], but see [22]). It should be noted, however, that the impact of channel expression level on TRPC6 regulation has not yet been addressed.

Evidently, despite some evidence that, at least under certain conditions some TRP proteins may form, or be part of store-operated Ca²⁺-permeable channels, the role of TRPs as genuine SOCs is still under debate. A chronic problem in recognizing TRPs as SOCs is the inability of ectopically expressed TRP proteins to recapitulate I_{CRAC}. However, it should be noted that although I_{CRAC} was the first store-operated Ca²⁺ current described and thus stands as the archetypical one, SOCs are heterogeneous with respect to both cation selectivity and pharmacological profile in different cell types (see [86] and refs. therein). This suggests that SOC channels may comprise a heterogeneous group of Ca²⁺-permeable channel proteins. Therefore, it is reasonable to conceive that SOC currents with different properties from I_{CRAC} may indeed involve channels built of TRP proteins. Within this context, it is important to note that functional TRPC channels are thought to be formed by four TRPC proteins that form either homo- or hetero-tetramers (see [140] and references therein) and, regardless of limitations as to which TRPCs can come together to form complete channels, it is recognized that the ultimate tetramer composition can alter functional and pharmacological features of TRPC channels [58,60,124]. Consequently, it is likely that native SOCs result from combinations of endogenous proteins not readily recapitulated in overexpression experiments. Undoubtedly, careful examination of the relationship between TRP protein expression level and channel regulation, as well as more emphasis on determining the composition of native TRP channels by taking advantage of knock out models and/or siRNA strategies, might eventually clarify existing discrepancies that hinder definition of their mode of regulation. However, even in those instances where TRP proteins have been found to be genuinely store-operated, the underlying mechanism coupling store-depletion to channel activation remains unknown. Within that context, in view of the recent discovery that the mammalian Ca²⁺-binding protein STIM1 (see Section 4.1 above) plays an essential role in the activation of SOCs, likely by acting as the sensor of Ca²⁺ store content [59,110], upcoming studies should address the potential role of STIM1 in regulation of store-operated TRPs.

6. Role of store-operated entry in [Ca²⁺]_i Oscillations

[Ca²⁺]_i oscillations, which generally consist of a series of repetitive spikes of [Ca²⁺]_i, represent the physiological response

of many cells to a wide variety of stimuli [4,6,129,130,148]. Agonist-induced oscillations usually occur as a result of PLC activation followed by the production of IP₃ and the repetitive release of Ca²⁺ from internal stores. At high agonist concentrations, it is widely accepted that SOCs are responsible for refilling the ER following store depletion [86,98]. However, the particular role that SOCs play in [Ca²⁺]_i oscillations, which occur at submaximal agonist concentrations and involve only a modest depletion of stores, is still a matter of debate.

Since the behavior of Ca²⁺ oscillations varies among different cell types as well as the type of agonist used, it is often difficult to resolve the complex relationship between oscillations and pool depletion. In many cell types, oscillations are abolished when Ca²⁺ is removed from the extracellular medium, suggesting that Ca²⁺ entry across the plasma membrane is necessary to maintain them [12,109,129,149]. It has been proposed that this may be achieved through the action of store-operated entry [5,129]. Conversely, Shuttleworth has provided evidence that agonist-induced oscillations are not mediated by SOCs, but involve Ca²⁺ entry channels regulated by lipid second messengers derived from products of PLC or PLA₂ [116]. Others have shown that other non-store operated channels such as TRPs are involved in the maintenance of oscillations [40,108,133,151].

Elimination of [Ca²⁺]_i oscillations in Ca²⁺-free external solutions suggests a role for Ca²⁺ entry in maintaining Ca²⁺ oscillations, but does not provide information on the precise nature of the Ca²⁺ entry pathway. In light of this, there is pharmacological evidence to support the involvement of store-operated entry. In one study, hormone-induced oscillations in hepatocytes were inhibited by Gd³⁺, 2-APB, and SK & F 96365, all at concentrations that inhibit SOCs and, specifically, I_{CRAC} [38,39]. In this same study, LOE-908, which blocks non-store operated channels, did not affect oscillations. Others have shown that La³⁺, another inhibitor of SOCs, blocks agonist-induced oscillations in cortical astrocytes [94]. In other studies using similar pharmacological tools, there is evidence that SOCs mediate oscillations in lymphocytes, HEK293 cells, and myotubes [12,28,55]. Whether entry through SOCs initiates or maintains IP₃-mediated Ca²⁺ oscillations remains unclear, especially since the molecular identity of the SOCs involved is unknown.

Despite the growing body of evidence to support the role that SOCs play in the maintenance of [Ca²⁺]_i oscillations, arguments have been raised against their involvement based on two observations. Shuttleworth and colleagues [116,117] maintain that agonist-induced oscillations are predominantly dependent on the generation of AA, a pathway that is activated independently of store depletion. In reports using HEK293 cells that overexpress the M3 muscarinic receptor and pancreatic acinar cells, addition of isotetrandrine, a phospholipase A2 blocker, inhibited the carbachol-induced oscillatory response [74,75]. Moreover, Shuttleworth and colleagues report that with high agonist concentrations, ARC channel activity is lost and the mode of Ca²⁺ entry occurs primarily through SOC activation, a phenomenon termed “reciprocal regulation” [74]. A similar reciprocal relationship between arachidonic acid-

activated and store-operated entry was described by Luo et al. [62] and by Moneer et al. [76].

TRP channels, which are associated with both SOC and ROC entry, have also been implicated in the maintenance of oscillations. One study identified a role for DAG-sensitive TRPC3 activation in the initiation of oscillations in glial cells [40]. OAG (1-oleoyl-2-acetyl-*sn*-glycerol; a synthetic DAG analog) -induced oscillations were preserved when external Ca²⁺ was replaced with Sr²⁺, providing further evidence for TRPC channel involvement since these channels are more permeable to Sr²⁺ cations than are SOCs. In another study using antisense technology to silence endogenous TRPC4, carbachol-induced oscillations were damped in HEK-293 cells in a manner similar to the run-down of oscillations in wild type HEK293 cells in Ca²⁺-free conditions [151].

The information encoded by [Ca²⁺]_i oscillations has been shown to control a variety of processes including gene transcription, muscle contraction, and differentiation. Hence, it is of great importance to elucidate the mechanisms underlying their initiation and maintenance. The Ca²⁺ entry during IP₃-mediated oscillations is difficult to resolve since the release from stores is small. However, in many cell types, oscillations abruptly cease or gradually run down when Ca²⁺ is removed from the external environment, making it clear that entry from the outside is a necessary component in this type of signaling process [42]. With the recent findings that STIM1 and Orai1 play an essential role in I_{CRAC} regulation (see above), these gene products may provide further clues as to the regulation of agonist-induced oscillations.

7. Pathophysiology associated with store-operated entry

Channelopathies are commonly the direct cause of disease, or contribute indirectly to the development of a disease. Knowing the extreme importance of SOC and CRAC channel activities in a variety of physiological functions, one would expect to see a pathophysiological phenotype if impaired store-operated entry occurred. The only documented disease state directly associated with a dysfunction in store-operated entry is a hereditary form of SCID syndrome [30,32,33]. SCID refers to a family of pathologies characterized by a malfunction in T and or B-lymphocyte activation and may be caused by a number of molecular abnormalities including mutations in cell surface receptors, signal transduction molecules, and transcription factors [27]. Regardless of the etiology of the condition, SCID patients share in common a hypersensitivity to infection due to compromised immune function. The form of SCID that is related to a dysfunction of store-operated entry was first attributed to a defect in nuclear import of the transcription factor NFAT in T-lymphocytes derived from affected patients [30]. In T-lymphocytes from normal individuals, presentation of an antigen causes a series of synchronized events, including the production of IP₃ and release of Ca²⁺ from the ER. This release of Ca²⁺ from the internal stores triggers influx of Ca²⁺ through CRAC channels, and this Ca²⁺ influx activates, via calmodulin, the phosphatase calcineurin which in turn dephosphorylates NFAT and induces nuclear import of the transcription factor

[31,67]. It was subsequently shown that the T-lymphocytes from patients suffering from this hereditary form of SCID completely lack I_{CRAC} [32]. Further, it was recently shown using a combination of two unbiased genome-wide approaches that a missense mutation, whereby a conserved arginine residue is replaced with a tryptophan at position 91 of the Orai1 gene, is responsible for the defect in I_{CRAC} in these patients [33]. Thus, not only has this series of studies with SCID patients led to the discovery of the molecular basis of a rare and debilitating disease, it has also proven instrumental in the discovery of a critical factor, Orai1, involved in the regulation of I_{CRAC} .

There is also evidence for a role of store-operated entry in diverse disease states such as acute pancreatitis [104] and Alzheimer's disease [70,72,154], as well as a possible role in causing some of the toxic effects of environmental chemicals that alter Ca^{2+} homeostasis [86]. Furthermore, there are to date four known channelopathies directly associated with TRP channels [79]. Three of these channelopathies involve TRPM6, TRPP2, and TRPML1, which have no known functions in store-operated entry. However, of particular interest to this discussion is the finding that TRPC6 is essential in the regulation of renal podocyte structure and function, and that a mutation in this channel that results in larger TRPC6 currents leads to proteinuria followed by progressive kidney failure [105,147]. However, as stated earlier, there is as yet no evidence to indicate that TRPC6 can function as a SOC under any conditions; therefore, it is not clear whether this defect in TRPC6 reflects a dysfunction in store-operated entry.

8. Conclusions

It has been 20 years since the commencement of study of store-operated Ca^{2+} entry pathways. And in that time, an extensive literature addressing properties of various SOC currents, pharmacological and cell biological properties of SOCs, and physiological and pathophysiological roles for store-operated entry has developed. However, until the very recent identification of roles for STIM1 and Orai1 in physiological I_{CRAC} , the field has suffered from a lack of unequivocal identification of molecular components of store-operated entry pathways. Thus, these are exciting times for the field of store-operated Ca^{2+} influx. The immediate challenge is of course to precisely define the mechanisms by which STIM1 and Orai1 regulate I_{CRAC} , as well as possibly other store-dependent entry pathways (Fig. 1). But beyond that, we are also faced with the task of revisiting much of this literature that has accumulated over the past 20 years in order to determine how the known functions of STIM1 and Orai1 may help address some of the outstanding, unresolved issues. For example, does STIM1 function as a universal Ca^{2+} sensor, capable of activating mechanisms of store-operated entry other than I_{CRAC} ? And if Orai1 is indeed the CRAC channel, then how do molecular determinants within the protein sequence explain the electrophysiological properties of I_{CRAC} such as permeation and ionic selectivity? Answers to these questions and many more will significantly enhance our understanding




Fig. 1. Proposed model of regulation of store-operated Ca^{2+} entry by STIM1 and Orai1. Agonist (Ag) stimulation of plasma membrane receptors (R) induces PLC activity, the products of which can directly activate TRPC channels in a receptor-operated fashion. Further, IP_3 generated by PLC induces release of Ca^{2+} from the ER via the IP_3R . STIM1, located within the ER membrane, senses ER Ca^{2+} depletion and translocates toward the plasma membrane, where it interacts with and activates Orai1, the putative CRAC channel. It remains to be determined whether STIM1 also regulates other store-dependent entry pathways, such as those mediated by store-operated TRPC channels.

of cellular Ca^{2+} signaling and its contribution to normal human health and disease.

References

- [1] J.M. Alderton, S.A. Ahmed, L.A. Smith, R.A. Steinhardt, Evidence for a vesicle-mediated maintenance of store-operated calcium channels in a human embryonic kidney cell line, *Cell Calcium* 28 (2000) 161.
- [2] D. Bakowski, M.D. Glitsch, A.B. Parekh, An examination of the secretion-like coupling model for the activation of the Ca^{2+} release-activated Ca^{2+} current I_{crac} in RBL-1 cells, *J. Physiol.* 532 (2001) 55 (Lond.).
- [3] D.J. Beech, S.Z. Xu, D. McHugh, R. Flemming, TRPC1 store-operated cationic channel subunit, *Cell Calcium* 33 (2003) 433.
- [4] M.J. Berridge, Calcium oscillations, *J. Biol. Chem.* 265 (1990) 9583.
- [5] M.J. Berridge, Cytoplasmic calcium oscillations: a two pool model, *Cell Calcium* 12 (1991) 63.
- [6] M.J. Berridge, Inositol trisphosphate and calcium oscillations, *Adv. Second Messenger Phosphoprot. Res.* 26 (1992) 211.
- [7] M.J. Berridge, Inositol trisphosphate and calcium signalling, *Nature* 361 (1993) 315.
- [8] M.J. Berridge, Capacitative calcium entry, *Biochem J.* 312 (1995) 1.
- [9] M.J. Berridge, P. Lipp, M.D. Bootman, The versatility and universality of calcium signalling, *Nat. Rev., Mol. Cell Biol.* 1 (2000) 11.
- [10] V.J. Bezzerides, I.S. Ramsey, S. Kotecha, A. Greka, D.E. Clapham, Rapid vesicular translocation and insertion of TRP channels, *Nat. Cell Biol.* 6 (2004) 709.
- [11] G.S. Bird, J.P. O.Aziz, B.J. Lievreumont, M. Wedel, G. Trebak, J. Vazquez Jr., Mechanisms of phospholipase C-regulated calcium entry, *Curr. Mol. Med.* 4 (2004) 291.
- [12] G.S. Bird, J. W.Putney Jr., Capacitative calcium entry supports calcium oscillations in human embryonic kidney cells, *J. Physiol.* 562 (2005) 697.
- [13] L. Bimbaumer, X. Zhu, M. Jiang, G. Boulay, M. Peyton, B. Vannier, D. Brown, D. Platano, H. Sadeghi, E. Stefani, M. Birnbaumer, On the molecular basis and regulation of cellular capacitative calcium entry: roles for Trp proteins, *Proc. Natl. Acad. Sci. U. S. A.* 93 (1996) 15195.
- [14] V.M. Bolotina, P. Csutora, CIF and other mysteries of the store-operated Ca^{2+} -entry pathway, *Trends Biochem. Sci.* 30 (2005) 378.
- [15] F.-J. Braun, L.M. Broad, D.L. Armstrong, J. W.Putney Jr., Stable activation of single CRAC-channels in divalent cation-free solutions, *J. Biol. Chem.* 276 (2001) 1063.

- [16] L. M. Broad, F.-J. Braun, J.-P. Lièvreumont, G. St. J. Bird, T. Kurosaki, J. W. Putney Jr., Role of the phospholipase C-inositol 1,4,5-trisphosphate pathway in calcium release-activated calcium current (I_{crac}) and capacitative calcium entry, *J. Biol. Chem.* 276 (2001) 15945.
- [17] L.M. Broad, T.R. Cannon, C.W. Taylor, A non-capacitative pathway activated by arachidonic acid is the major Ca²⁺ entry mechanism in rat A7r5 smooth muscle cells stimulated with low concentrations of vasopressin, *J. Physiol. (London)* 517 (1999) 121.
- [18] G.H. Brough, S. Wu, D. Cioffi, T.M. Moore, M. Li, N. Dean, T. Stevens, Contribution of endogenously expressed Trp1 to a Ca²⁺-selective, store-operated Ca²⁺ entry pathway, *FASEB J.* 15 (2001) 1727.
- [19] L.I. Brueggemann, D.R. Markun, K.K. Henderson, L.L. Cribbs, K.L. Byron, Pharmacological and electrophysiological characterization of store-operated currents and capacitative Ca(2+) entry in vascular smooth muscle cells, *J Pharmacol. Exp. Ther.* 317 (2006) 488.
- [20] D. Burdakov, O.H. Petersen, A. Verkhratsky, Intraluminal calcium as a primary regulator of endoplasmic reticulum function, *Cell Calcium* 38 (2005) 303.
- [21] C. Camello, R. Lomax, O.H. Petersen, A. Tepikin, Calcium leak from intracellular stores—The enigma of calcium signalling, *Cell Calcium* 32 (2002) 355.
- [22] S. Cayouette, M.P. Lussier, E.L. Mathieu, S.M. Bousquet, G. Boulay, Exocytotic insertion of TRPC6 channel into the plasma membrane upon Gq protein-coupled receptor activation, *J. Biol. Chem.* 279 (2004) 7241.
- [23] M.K. Chung, H. Lee, A. Mizuno, M. Suzuki, M.J. Caterina, 2-aminoethoxydiphenyl borate activates and sensitizes the heat-gated ion channel TRPV3, *J. Neurosci.* 24 (2004) 5177.
- [24] E. Clementi, A. Martini, G. Stefani, J. Meldolesi, P. Volpe, LU52396, an inhibitor of the store-dependent (capacitative) Ca²⁺ influx, *Eur. J. Pharmacol. Mol. Pharmacol.* 289 (1995) 23.
- [25] E. Clementi, H. Scheer, D. Zacchetti, C. Fasolato, T. Pozzan, J. Meldolesi, Receptor-activated Ca²⁺ influx. Two independently regulated mechanisms of influx stimulation coexist in neurosecretory PC12 cells, *J. Biol. Chem.* 267 (1992) 2164.
- [26] P. Csutora, Z. Su, H.Y. Kim, A. Bugrim, K.W. Cunningham, R. Nuccitelli, J.E. Keizer, M.R. Hanley, J.E. Blalock, R.B. Marchase, Calcium influx factor is synthesized by yeast and mammalian cells depleted of organellar calcium stores, *Proc. Natl. Acad. Sci. U. S. A.* 96 (1999) 121.
- [27] C. Cunningham-Rundles, P.P. Ponda, Molecular defects in T- and B-cell primary immunodeficiency diseases, *Nat. Rev., Immunol.* 5 (2005) 880.
- [28] M. Estrada, A. Espinosa, C.J. Gibson, P. Uhlen, E. Jaimovich, Capacitative calcium entry in testosterone-induced intracellular calcium oscillations in myotubes, *J. Endocrinol.* 184 (2005) 371.
- [29] C.C. Felder, A.L. Ma, L.A. Liotta, E.C. Kohn, The antiproliferative and antimetastatic compound L615182, inhibits muscarinic acetylcholine receptor-stimulated calcium influx and arachidonic acid release, *J. Pharmacol. Exp. Ther.* 257 (1991) 967.
- [30] S. Feske, R. Draeger, H.H. Peter, A. Rao, Impaired NFAT regulation and its role in a severe combined immunodeficiency, *Immunobiology* 202 (2000) 134.
- [31] S. Feske, H. Okamura, P.G. Hogan, A. Rao, Ca²⁺/calcineurin signalling in cells of the immune system, *Biochem. Biophys. Res Commun.* 311 (2003) 1117.
- [32] S. Feske, M. Prakriya, A. Rao, R.S. Lewis, A severe defect in CRAC Ca²⁺ channel activation and altered K⁺ channel gating in T cells from immunodeficient patients, *J. Exp. Med.* 202 (2005) 651.
- [33] S. Feske, Y. Gwack, M. Prakriya, S. Srikanth, S.H. Puppel, B. Tanasa, P.G. Hogan, R.S. Lewis, M. Daly, and A. Rao, A mutation in Orail causes immune deficiency by abrogating CRAC channel function, *Nature* 441 (2006) 179.
- [34] D. Franzius, M. Hoth, R. Penner, Non-specific effects of calcium entry antagonists in mast cells, *Pflügers Arch.* 428 (1994) 433.
- [35] M. Freichel, S.H. Suh, A. Pfeifer, U. Schweig, C. Trost, P. Weißgerber, M. Biel, S. Philipp, D. Freise, G. Droogmans, F. Hofmann, V. Flockerzi, B. Nilius, Lack of an endothelial store-operated Ca²⁺ current impairs agonist-dependent vasorelaxation in TRP4^{-/-} mice, *Nat. Cell Biol.* 3 (2001) 121.
- [36] M. Freichel, R. Vennekens, J. Olausson, S. Stolz, S.E. Philipp, P. Weissgerber, V. Flockerzi, Functional role of TRPC proteins in native systems: implications from knockout and knock-down studies, *J. Physiol.* 567 (2005) 59.
- [37] P. Gailly, M. Colson-Van Schoor, Involvement of trp-2 protein in store-operated influx of calcium in fibroblasts, *Cell Calcium* 30 (2001) 157.
- [38] R.B. Gregory, G.J. Barritt, Evidence that Ca²⁺-release-activated Ca²⁺ channels in rat hepatocytes are required for the maintenance of hormone-induced Ca²⁺ oscillations, *Biochem. J.* 370 (2003) 695.
- [39] R.B. Gregory, G. Rychkov, G.J. Barritt, Evidence that 2-aminoethyl diphenylborate is a novel inhibitor of store-operated Ca²⁺ channels in liver cells, and acts through a mechanism which does not involve inositol trisphosphate receptors, *Biochem. J.* 354 (2001) 285.
- [40] M. Grimaldi, M. Maratos, A. Verma, Transient receptor potential channel activation causes a novel form of [Ca²⁺] oscillations and is not involved in capacitative Ca²⁺ entry in glial cells, *J Neurosci.* 23 (2003) 4737.
- [41] R.W. Gross, Activation of calcium-independent phospholipase A2 by depletion of internal calcium stores, *Biochem. Soc. Trans.* 26 (1998) 345.
- [42] S.D. Hess, M. Oortgiesen, M.D. Cahalan, Calcium oscillations in human T and natural killer cells depend upon membrane potential and calcium influx, *J. Immunol.* 150 (1993) 2620.
- [43] C. Hisatsune, K. Mikoshiba, Novel compartment implicated in calcium signaling—Is it an “induced coupling domain”? *Sci STKE* 2005 (2005) e53.
- [44] T. Hofmann, M. Schaefer, G. Schultz, T. Gudermann, Cloning, expression and subcellular localization of two splice variants of mouse transient receptor potential channel 2, *Biochem. J.* 351 (2000) 115.
- [45] M. Hoth, R. Penner, Depletion of intracellular calcium stores activates a calcium current in mast cells, *Nature* 355 (1992) 353.
- [46] M. Hoth, R. Penner, Calcium release-activated calcium current in rat mast cells, *J. Physiol. (London)* 465 (1993) 359.
- [47] H.Z. Hu, Q. Gu, C. Wang, C.K. Colton, J. Tang, M. Kinoshita-Kawada, L.Y. Lee, J.D. Wood, M.X. Zhu, 2-aminoethoxydiphenyl borate is a common activator of TRPV1, TRPV2, and TRPV3, *J. Biol. Chem.* 279 (2004) 35741.
- [48] R.F. Irvine, “Quantal” Ca²⁺ release and the control of Ca²⁺ entry by inositol phosphates—A possible mechanism, *FEBS Lett.* 263 (1990) 5.
- [49] M.K. Jungnickel, H. Marreo, L. Birnbaumer, J.R. Lemos, H.M. Florman, Trp2 regulates entry of Ca²⁺ into mouse sperm triggered by egg ZP3, *Nat. Cell Biol.* 3 (2001) 499.
- [50] H.Y. Kim, D. Thomas, M.R. Hanley, Chromatographic resolution of an intracellular calcium influx factor from thapsigargin-activated Jurkat cells, *J. Biol. Chem.* 270 (1995) 9706.
- [51] K. Kiselyov, G.A. Mignery, M.X. Zhu, S. Muallem, The N-terminal domain of the IP3 receptor gates store-operated hTrp3 channels, *Mol. Cell* 4 (1999) 423.
- [52] K. Kiselyov, X. Xu, G. Mozhayeva, T. Kuo, I. Pessah, G. Mignery, X. Zhu, L. Birnbaumer, S. Muallem, Functional interaction between InsP3 receptors and store-operated Htrp3 channels, *Nature* 396 (1998) 478.
- [53] Y.-M. Leung, C.-Y. Kwan, T.-T. Loh, Dual effects of tetrandrine on cytosolic calcium in human leukaemic HL-60 cells: intracellular calcium release and calcium entry blockade, *Br. J. Pharmacol.* 113 (1994) 767.
- [54] R.S. Lewis, Calcium signaling mechanisms in T lymphocytes, *Annu. Rev. Immunol.* 19 (2001) 497.
- [55] R.S. Lewis, Calcium oscillations in T-cells: mechanisms and consequences for gene expression, *Biochem. Soc. Trans.* 31 (2003) 925.
- [56] J.P. Lievreumont, G.S. Bird, J.W. Putney Jr., Canonical transient receptor potential TRPC7 can function as both a receptor- and store-operated channel in HEK-293 cells, *Am. J. Physiol.: Cell Physiol.* 287 (2004) C1709–C1716.
- [57] J.P. Lievreumont, T. Numaga, G. Vazquez, L. Lemonnier, Y. Hara, E. Mori, M. Trebak, S.E. Moss, G.S. Bird, Y. Mori, J.W. Putney Jr., The Role of Canonical Transient Receptor Potential 7 in B-cell Receptor-activated Channels, *J. Biol. Chem.* 280 (2005) 35346.
- [58] B. Lintschinger, M. Balzer-Geldsetzer, T. Baskaran, W.F. Graier, C. Romanin, M.X. Zhu, K. Groschner, Coassembly of Trp1 and Trp3 proteins generates diacylglycerol- and Ca²⁺-sensitive cation channels, *J. Biol. Chem.* 275 (2000) 27799.

- [59] J. Liou, M.L. Kim, W.D. Heo, J.T. Jones, J.W. Myers Jr., J.E. Ferrell, Jr., and T. Meyer, STIM is a Ca²⁺ sensor essential for Ca²⁺-store-depletion-triggered Ca²⁺ influx, *Curr. Biol.* 15 (2005) 1235.
- [60] X. Liu, B.C. Bandyopadhyay, B.B. Singh, K. Groschner, I.S. Ambudkar, Molecular analysis of a store-operated and 2-acetyl-sn-glycerol-sensitive non-selective cation channel. Heteromeric assembly of TRPC1–TRPC3, *J. Biol. Chem.* 280 (2005) 21600.
- [61] X. Liu, W. Wang, B.B. Singh, T. Lockwich, J. Jadowiec, B. O'Connell, R. Wellner, M.X. Zhu, I.S. Ambudkar, Trp1, a candidate protein for the store-operated Ca²⁺ influx mechanism in salivary gland cells, *J. Biol. Chem.* 275 (2000) 3403.
- [62] D. Luo, L.M. Broad, G.S. Bird, J.W. Putney Jr., Mutual antagonism of calcium entry by capacitative and arachidonic acid-mediated calcium entry pathways, *J. Biol. Chem.* 276 (2001) 20186.
- [63] D. Luo, L.M. Broad, G.St.J. Bird, J.W. Putney Jr., Signaling pathways underlying muscarinic receptor-induced [Ca²⁺]_i oscillations in HEK293 cells, *J. Biol. Chem.* 276 (2001) 5613.
- [64] H.-T. Ma, R.L. Patterson, D.B. van Rossum, L. Birnbaumer, K. Mikoshiba, D.L. Gill, Requirement of the inositol trisphosphate receptor for activation of store-operated Ca²⁺ channels, *Science* 287 (2000) 1647.
- [65] H.-T. Ma, K. Venkatachalam, H.-S. Li, C. Montell, T. Kuroski, R.L. Patterson, D.L. Gill, Assessment of the role of the inositol 1,4,5-trisphosphate receptor in the activation of transient receptor potential channels and store-operated Ca²⁺ entry channels, *J. Biol. Chem.* 276 (2001) 18888.
- [66] H.-T. Ma, K. Venkatachalam, J.B. Parys, D.L. Gill, Modification of store-operated channel coupling and inositol trisphosphate receptor function by 2-aminoethoxydiphenyl borate in DT40 lymphocytes, *J. Biol. Chem.* 277 (2002) 6915.
- [67] F. Macian, NFAT proteins: key regulators of T-cell development and function, *Nat. Rev. Immunol.* 5 (2005) 472.
- [68] S.S. Manji, N.J. Parker, R.T. Williams, S.L. Van, R.B. Pearson, M. Dziadek, P.J. Smith, STIM1: a novel phosphoprotein located at the cell surface, *Biochim. Biophys. Acta* 1481 (2000) 147.
- [69] T. Maruyama, T. Kanaji, S. Nakade, T. Kanno, K. Mikoshiba, 2APB, 2-aminoethoxydiphenyl borate, a membrane-penetrable modulator of Ins (1,4,5)P₃-induced Ca²⁺ release, *J. Biochem.* 122 (1997) 498.
- [70] M.P. Mattson, S.L. Chan, Neuronal and glial calcium signaling in Alzheimer's disease, *Cell Calcium* 34 (2003) 385.
- [71] R.R. McKay, C.L. Szmeczek-Seay, J.-P. Lièvreumont, G.St.J. Bird, C. Zitt, E. Jüngling, A. Lückhoff, J.W. Putney Jr., Cloning and expression of the human transient receptor potential 4 (TRP4) gene: localization and functional expression of human TRP4 and TRP3, *Biochem. J.* 351 (2000) 735.
- [72] J.G. McLarnon, H.B. Choi, L.F. Lue, D.G. Walker, S.U. Kim, Perturbations in calcium-mediated signal transduction in microglia from Alzheimer's disease patients, *J. Neurosci. Res.* 81 (2005) 426.
- [73] J.C. Mercer, W.I. DeHaven, J.T. Smyth, B. Wedel, R.R. Boyles, G.S. Bird, J.W. Putney Jr., Large store-operated calcium-selected currents due to co-expression of orai1 or orai2 with the intracellular calcium sensor, stim1, *J. Biol. Chem.* 281 (2006) 24979.
- [74] O. Mignen, J.L. Thompson, T.J. Shuttleworth, Reciprocal regulation of capacitative and arachidonate-regulated noncapacitative Ca²⁺ entry pathways, *J. Biol. Chem.* 276 (2001) 35676.
- [75] O. Mignen, J.L. Thompson, D.I. Yule, T.J. Shuttleworth, Agonist activation of arachidonate-regulated Ca²⁺-selective (ARC) channels in murine parotid and pancreatic acinar cells, *J. Physiol.* 564 (2005) 791.
- [76] Z. Moneer, C.W. Taylor, Reciprocal regulation of capacitative and non-capacitative Ca²⁺ entry in A7r5 vascular smooth muscle cells: only the latter operates during receptor activation, *Biochem. J.* 362 (2002) 12.
- [77] C. Montell, TRP channels in *Drosophila* photoreceptor cells, *J. Physiol.* 567 (2005) 45.
- [78] Y. Mori, M. Wakamori, T. Miyakawa, M. Hermosura, Y. Hara, M. Nishida, K. Hirose, A. Mizushima, M. Kuroski, E. Mori, K. Gotoh, T. Okada, A. Fleig, R. Penner, M. Iino, T. Kuroski, Transient receptor potential 1 regulates capacitative Ca²⁺ entry and Ca²⁺ release from endoplasmic reticulum in B lymphocytes, *J. Exp. Med.* 195 (2002) 673.
- [79] B. Nilius, T. Voets, J. Peters, TRP channels in disease, *Sci. STKE* 2005 (2005) re8.
- [80] A.G. Obukhov, M.C. Nowycky, TRPC4 can be activated by G-protein-coupled receptors and provides sufficient Ca²⁺ to trigger exocytosis in neuroendocrine cells, *J. Biol. Chem.* 277 (2002) 16172.
- [81] A.F. Odell, J.L. Scott, D.F. Van Helden, Epidermal Growth Factor Induces Tyrosine Phosphorylation, Membrane Insertion, and Activation of Transient Receptor Potential Channel 4, *J. Biol. Chem.* 280 (2005) 37974.
- [82] T. Okada, S. Shimizu, M. Wakamori, A. Maeda, T. Kuroski, N. Takada, K. Imoto, Y. Mori, Molecular cloning and functional characterization of a novel receptor-activated TRP Ca²⁺ channel from mouse brain, *J. Biol. Chem.* 273 (1998) 10279.
- [83] T. Okada, R. Inoue, K. Yamazaki, A. Maeda, T. Kuroski, T. Yamakuni, I. Tanaka, S. Shimizu, K. Ikenaka, K. Imoto, Y. Mori, Molecular and functional characterization of a novel mouse transient receptor potential protein homologue TRP7. Ca²⁺-permeable cation channel that is constitutively activated and enhanced by stimulation of G protein-coupled receptor, *J. Biol. Chem.* 274 (1999) 27359.
- [84] B. Ordaz, J. Tang, R. Xiao, A. Salgado, A. Sampieri, M.X. Zhu, L. Vaca, Calmodulin and calcium interplay in the modulation of TRPC5 channel activity. Identification of a novel C-terminal domain for calcium/calmodulin-mediated facilitation, *J. Biol. Chem.* 280 (2005) 30788.
- [85] A.B. Parekh, R. Penner, Store depletion and calcium influx, *Physiol. Rev.* 77 (1997) 901.
- [86] A.B. Parekh, J.W. Putney Jr., Store-operated calcium channels, *Physiol. Rev.* 85 (2005) 757.
- [87] R.L. Patterson, D.B. van Rossum, D.L. Ford, K.J. Hurt, S.S. Bae, P.-G. Suh, T. Kuroski, S.H. Snyder, D.L. Gill, Phospholipase C-g is required for agonist-induced Ca²⁺ entry, *Cell* 111 (2002) 529.
- [88] R.L. Patterson, D.B. van Rossum, D.L. Gill, Store-operated Ca²⁺ entry: evidence for a secretion-like coupling model, *Cell* 98 (1999) 487.
- [89] S.F. Pedersen, G. Owsianik, B. Nilius, TRP channels: an overview, *Cell Calcium* 38 (2005) 233.
- [90] C. Peinelt, M. Vig, D.L. Koomoa, A. Beck, M.J. Nadler, M. Koblan-Huberson, A. Lis, A. Fleig, R. Penner, J.P. Kinet, Amplification of CRAC current by STIM1 and CRACM1 (Orai1), *Nat. Cell Biol.* 8 (2006) 771.
- [91] S. Philipp, A. Cavalié, M. Freichel, U. Wissenbach, S. Zimmer, C. Trost, A. Marguaret, M. Murakami, V. Flockerzi, A mammalian capacitative calcium entry channel homologous to *Drosophila* TRP and TRPL, *EMBO J.* 15 (1996) 6166.
- [92] S. Philipp, J. Hambrecht, L. Braslavski, G. Schroth, M. Freichel, M. Murakami, A. Cavalié, V. Flockerzi, A novel capacitative calcium entry channel expressed in excitable cells, *EMBO J.* 17 (1998) 4274.
- [93] S. Philipp, C. Trost, J. Warnat, J. Rautmann, N. Himmerkus, G. Schroth, O. Kretz, W. Nastainczyk, A. Cavalié, M. Hoth, V. Flockerzi, Trp4 (CCE1) protein is part of native calcium release-activated Ca²⁺-like channels in adrenal cells, *J. Biol. Chem.* 275 (2000) 23965.
- [94] P. Pizzo, A. Burgo, T. Pozzan, C. Fasolato, Role of capacitative calcium entry on glutamate-induced calcium influx in type-I rat cortical astrocytes, *J. Neurochem.* 79 (2001) 98.
- [95] T.D. Plant, M. Schaefer, TRPC4 and TRPC5: receptor-operated Ca²⁺-permeable nonselective cation channels, *Cell Calcium* 33 (2003) 441.
- [96] M. Prakriya, R.S. Lewis, Potentiation and inhibition of Ca²⁺ release-activated Ca²⁺ channels by 2-aminoethoxydiphenyl borate (2-APB) occurs independently of IP₃ receptors, *J. Physiol.* 536 (2001) 3 (Lond.).
- [97] M. Prakriya, R.S. Lewis, CRAC channels: activation, permeation, and the search for a molecular identity, *Cell Calcium* 33 (2003) 311.
- [98] J.W. Putney Jr., A model for receptor-regulated calcium entry, *Cell Calcium* 7 (1986) 1.
- [99] J.W. Putney Jr., Pharmacology of capacitative calcium entry, *Mol. Interv.* 1 (2001) 84.
- [100] J.W. Putney Jr., Store-operated calcium channels: how do we measure them, and why do we care? *Sci. STKE* 2004 (2004) e37.
- [101] J.W. Putney Jr., The enigmatic TRPCs: multifunctional cation channels, *Trends Cell Biol.* 14 (2004) 282.

- [102] I.S. Ramsey, M. Delling, D.E. Clapham, An introduction to TRP channels, *Annu. Rev. Physiol.* 68 (2006) 619.
- [103] C. Randriamampita, R.Y. Tsien, Emptying of intracellular Ca^{2+} stores releases a novel small messenger that stimulates Ca^{2+} influx, *Nature* 364 (1993) 809.
- [104] M. Raraty, J. Ward, G. Erdemli, C. Vaillant, J.P. Neoptolemos, R. Sutton, O.H. Petersen, Calcium-dependent enzyme activation and vacuole formation in the apical granular region of pancreatic acinar cells, *Proc. Natl. Acad. Sci. U. S. A.* 97 (2000) 13126.
- [105] J. Reiser, K.R. Polu, C.C. Moller, P. Kenlan, M.M. Altintas, C. Wei, C. Faul, S. Herbert, I. Villegas, C. Vila-Casado, M. McGee, H. Sugimoto, D. Brown, R. Kalluri, P. Mundel, P.L. Smith, D.E. Clapham, M.R. Pollak, TRPC6 is a glomerular slit diaphragm-associated channel required for normal renal function, *Nat. Genet.* 37 (2005) 739.
- [106] C.M.P. Ribeiro, J. Reece, J.W. Putney Jr., Role of the cytoskeleton in calcium signaling in NIH 3T3 cells. An intact cytoskeleton is required for agonist-induced $[\text{Ca}^{2+}]_i$ signaling, but not for capacitative calcium entry, *J. Biol. Chem.* 272 (1997) 26555.
- [107] A. Riccio, C. Mattei, R.E. Kelsell, A.D. Medhurst, A.R. Calver, A.D. Randall, J.B. Davis, C.D. Benham, M.N. Pangalos, Cloning and functional expression of human short TRP7, a candidate protein for store-operated Ca^{2+} influx, *J. Biol. Chem.* 277 (2002) 12302.
- [108] M.W. Roe, J.F. Worley III, F. Qian, N. Tamarina, A.A. Mittal, F. Dralyuk, N.T. Blair, R.J. Mertz, L.H. Philipson, I.D. Dukes, Characterization of a Ca^{2+} release-activated nonselective cation current regulating membrane potential and $[\text{Ca}^{2+}]_i$ oscillations in transgenically derived β -cells, *J. Biol. Chem.* 273 (1998) 10402.
- [109] T.A. Rooney, E.J. Sass, A.P. Thomas, Characterization of cytosolic calcium oscillations induced by phenylephrine and vasopressin in single fura-2-loaded hepatocytes, *J. Biol. Chem.* 264 (1989) 17131.
- [110] J. Roos, P.J. DiGregorio, A.V. Yeromin, K. Ohlsen, M. Lioudyno, S. Zhang, O. Saffrina, J.A. Kozak, S.L. Wagner, M.D. Cahalan, G. Velicelebi, K.A. Stauderman, STIM1, an essential and conserved component of store-operated Ca^{2+} channel function, *J. Cell Biol.* 169 (2005) 435.
- [111] J.A. Rosado, S.O. Sage, A role for the actin cytoskeleton in the initiation and maintenance of store-mediated calcium entry in human platelets, *Trends Cardiovasc. Med.* 10 (2000) 327.
- [112] B.A. Rzigalinski, P.F. Blackmore, M.D. Rosenthal, Arachidonate mobilization is coupled to depletion of intracellular calcium stores and influx of extracellular calcium in differentiated U937 cells, *Biochim. Biophys. Acta* 1299 (1996) 342.
- [113] M. Schaefer, T.D. Plant, A.G. Obukhov, T. Hofmann, T. Gudermann, G. Schultz, Receptor-mediated regulation of the nonselective cation channels TRPC4 and TRPC5, *J. Biol. Chem.* 275 (2000) 17517.
- [114] C.C. Scott, W. Furuya, W.S. Trimble, S. Grinstein, Activation of store-operated Calcium Channels: ASSESSMENT OF THE ROLE OF SNARE-MEDIATED VESICULAR TRANSPORT, *J. Biol. Chem.* 278 (2003) 30534.
- [115] S. Shimizu, T. Yoshida, M. Wakamori, M. Ishii, T. Okada, M. Takahashi, M. Seto, K. Sakurada, Y. Kiuchi, Y. Mori, Ca^{2+} -calmodulin-dependent myosin light chain kinase is essential for activation of TRPC5 channels expressed in HEK293 cells, *J. Physiol.* 570 (2006) 219.
- [116] T.J. Shuttleworth, What drives calcium entry during $[\text{Ca}^{2+}]_i$ oscillations? —Challenging the capacitative model, *Cell Calcium* 25 (1999) 237.
- [117] T.J. Shuttleworth, O. Mignen, Calcium entry and the control of calcium oscillations, *Biochem. Soc. Trans.* 31 (2003) 916.
- [118] B.B. Singh, T.P. Lockwich, B.C. Bandyopadhyay, X. Liu, S. Bollimuntha, S.C. Brazer, C. Combs, S. Das, A.G. Leenders, Z.H. Sheng, M.A. Knepper, S.V. Ambudkar, I.S. Ambudkar, VAMP2-Dependent Exocytosis Regulates Plasma Membrane Insertion of TRPC3 Channels and Contributes to Agonist-Stimulated Ca^{2+} Influx, *Mol. Cell* 15 (2004) 635.
- [119] T. Smani, S.I. Zakharov, P. Csutora, E. Leno, E.S. Trepakova, V.M. Bolotina, A novel mechanism for the store-operated calcium influx pathway, *Nat. Cell Biol.* 6 (2004) 113.
- [120] T. Smani, S.I. Zakharov, E. Leno, P. Csutora, E.S. Trepakova, V.M. Bolotina, Ca^{2+} -independent phospholipase A2 is a novel determinant of store-operated Ca^{2+} entry, *J. Biol. Chem.* 278 (2003) 11909.
- [121] J.T. Smyth, L. Lemonnier, G. Vazquez, G.S. Bird, J.W. Putney Jr., Dissociation of regulated trafficking of TRPC3 channels to the plasma membrane from their activation by phospholipase C, *J. Biol. Chem.* 281 (2005) 11712.
- [122] J. Soboloff, M.A. Spassova, X.D. Tang, T. Hewavitharana, W. Xu, D.L. Gill, Orai1 and STIM reconstitute store-operated calcium channel function, *J. Biol. Chem.* 281 (2006) 20661.
- [123] M. Spassova, J. Soboloff, L.-P. He, W. Xu, M. Dziadek, D.L. Gill, STIM1 has a plasma membrane role in the activation of store-operated Ca^{2+} channels, *Proc. Natl. Acad. Sci. U. S. A.* 103 (2006) 4040.
- [124] C. Strübing, G. Krapivinsky, L. Krapivinsky, D.E. Clapham, TRPC1 and TRPC5 form a novel cation channel in mammalian brain, *Neuron* 29 (2001) 645.
- [125] H. Sugawara, M. Kurosaki, M. Takata, T. Kurosaki, Genetic evidence for involvement of type 1, type 2 and type 3 inositol 1,4,5-trisphosphate receptors in signal transduction through the B-cell antigen receptor, *EMBO J.* 16 (1997) 3078.
- [126] M. Sweeney, Y. Yu, O. Platoshyn, S. Zhang, S.S. McDaniel, J.X. J. Yuan, Inhibition of endogenous TRP1 decreases capacitative Ca^{2+} entry and attenuates pulmonary artery smooth muscle cell proliferation, *Am. J. Physiol.* 283 (2002) L144–L155.
- [127] H. Takemura, A.R. Hughes, O. Thastrup, J.W. Putney Jr., Activation of calcium entry by the tumor promoter, thapsigargin, in parotid acinar cells. Evidence that an intracellular calcium pool, and not an inositol phosphate, regulates calcium fluxes at the plasma membrane, *J. Biol. Chem.* 264 (1989) 12266.
- [128] H. Takemura, J.W. Putney Jr., Capacitative calcium entry in parotid acinar cells, *Biochem. J.* 258 (1989) 409.
- [129] A.P. Thomas, G.St.J. Bird, G. Hajnóczky, L.D. Robb-Gaspers, J.W. Putney Jr., Spatial and temporal aspects of cellular calcium signalling, *FASEB J.* 10 (1996) 1505.
- [130] A.P. Thomas, D.C. Renard, T.A. Rooney, Spatial organization of Ca^{2+} signalling and $\text{Ins}(1,4,5)\text{P}_3$ action, *Adv. Second Messenger Phosphoprot. Res.* 26 (1992) 225.
- [131] D. Thomas, M.R. Hanley, Evaluation of calcium influx factors from stimulated Jurkat T-lymphocytes by microinjection into *Xenopus* oocytes, *J. Biol. Chem.* 270 (1995) 6429.
- [132] Y. Tomita, S. Kaneko, M. Funayama, H. Kondo, M. Satoh, A. Akaike, Intracellular Ca^{2+} store-operated influx of Ca^{2+} through TRP-R, a rat homolog of TRP, expressed in *Xenopus* oocytes, *Neurosci. Lett.* 248 (1998) 195.
- [133] S. Torihashi, T. Fujimoto, C. Trost, S. Nakayama, Calcium oscillation linked to pacemaking of interstitial cells of Cajal, *J. Biol. Chem.* 277 (2002) 19191.
- [134] M. Trebak, G.St.J. Bird, R.R. McKay, L. Birnbaumer, J.W. Putney Jr., Signaling mechanism for receptor-activated TRPC3 channels, *J. Biol. Chem.* 278 (2003) 16244.
- [135] M. Trebak, G.St.J. Bird, R.R. McKay, J.W. Putney Jr., Comparison of human TRPC3 channels in receptor-activated and store-operated modes. Differential sensitivity to channel blockers suggests fundamental differences in channel composition, *J. Biol. Chem.* 277 (2002) 21617.
- [136] M. Trebak, G. Vazquez, G.St.J. Bird, J.W. Putney Jr., The TRPC3/6/7 subfamily of cation channels, *Cell Calcium* 33 (2003) 451.
- [137] E.S. Trepakova, M. Gericke, Y. Hirakawa, R.M. Weisbrod, R.A. Cohen, V.M. Bolotina, Properties of a native cation channel activated by Ca^{2+} store depletion in vascular smooth muscle cells, *J. Biol. Chem.* 276 (2001) 7782.
- [138] B. Vannier, M. Peyton, G. Boulay, D. Brown, N. Qin, M. Jiang, X. Zhu, L. Birnbaumer, Mouse *trp2*, the homologue of the human *trpc2* pseudogene, encodes mTrp2, a store depletion-activated capacitative Ca^{2+} channel, *Proc. Natl. Acad. Sci. U. S. A.* 96 (1999) 2060.
- [139] G. Vazquez, J.-P. Lièvreumont, G.St.J. Bird, J.W. Putney Jr., Human Trp3 forms both inositol trisphosphate receptor-dependent and receptor-independent store-operated cation channels in DT40 avian B-lymphocytes, *Proc. Natl. Acad. Sci. U. S. A.* 98 (2001) 11777.
- [140] G. Vazquez, B.J. Wedel, O. Aziz, M. Trebak, J.W. Putney Jr., The mammalian TRPC cation channels, *Biochim. Biophys. Acta* 1742 (2004) 21.

- [141] G. Vazquez, B.J. Wedel, G.St.J. Bird, S.K. Joseph, J.W. Putney Jr., An inositol 1,4,5-trisphosphate receptor-dependent cation entry pathway in DT40 B lymphocytes, *EMBO J.* 21 (2002) 4531.
- [142] G. Vazquez, B.J. Wedel, M. Trebak, G.St.J. Bird, J.W. Putney Jr., Expression level of TRPC3 channel determines its mechanism of activation, *J. Biol. Chem.* 278 (2003) 21649.
- [143] M. Vig, C. Peinelt, A. Beck, D.L. Koomoa, D. Rabah, M. Koblan-Huberson, S. Kraft, H. Turner, A. Fleig, R. Penner, J.P. Kinet, CRACM1 Is a Plasma Membrane Protein Essential for store-operated Ca²⁺ Entry, *Science* 312 (2006) 1220.
- [144] J. Wamat, S. Philipp, S. Zimmer, V. Flockerzi, A. Cavalié, Phenotype of a recombinant store-operated channel: highly selective permeation of Ca²⁺, *J. Physiol. (London)* 518 (1999) 631.
- [145] P.D. Wes, J. Chevesich, A. Jeromin, C. Rosenberg, G. Stetten, C. Montell, TRPC1, a human homolog of a *Drosophila* store-operated channel, *Proc. Natl. Acad. Sci. U. S. A.* 92 (1995) 9652.
- [146] R.T. Williams, P.V. Senior, S.L. Van, J.E. Layton, P.J. Smith, M.A. Dziadek, Stromal interaction molecule 1 (STIM1), a transmembrane protein with growth suppressor activity, contains an extracellular SAM domain modified by N-linked glycosylation, *Biochim. Biophys. Acta* 1596 (2002) 131.
- [147] M.P. Winn, P.J. Conlon, K.L. Lynn, M.K. Farrington, T. Creazzo, A.F. Hawkins, N. Daskalakis, S.Y. Kwan, S. Ebersviller, J.L. Burchette, M.A. Pericak-Vance, D.N. Howell, J.M. Vance, P.B. Rosenberg, A mutation in the TRPC6 cation channel causes familial focal segmental glomerulosclerosis, *Science* 308 (2005) 1801.
- [148] N.M. Woods, K.S. Cuthbertson, P.H. Cobbold, Repetitive transient rises in cytoplasmic free calcium in hormone-stimulated hepatocytes, *Nature* 319 (1986) 600.
- [149] N.M. Woods, C.J. Dixon, K.S.R. Cuthbertson, P.H. Cobbold, Modulation of free Ca oscillations in single hepatocytes by changes in extracellular K⁺, Na⁺ and Ca²⁺, *Cell Calcium* 11 (1990) 353.
- [150] X. Wu, G. Babnigg, M.L. Villereal, Functional significance of human trp1 and trp3 in store-operated Ca²⁺ entry in HEK-293 cells, *Am. J. Physiol.* 278 (2000) C526–C536.
- [151] X. Wu, G. Babnigg, T. Zagranichnaya, M.L. Villereal, The role of endogenous human Trp4 in regulating carbachol-induced calcium oscillations in HEK-293 cells, *J. Biol. Chem.* 277 (2002) 13597.
- [152] H. Yang, S. Mergler, X. Sun, Z. Wang, L. Lu, J.A. Bonanno, U. Pleyer, P.S. Reinach, TRPC4 Knockdown Suppresses Epidermal Growth Factor-induced store-operated Channel Activation and Growth in Human Corneal Epithelial Cells, *J. Biol. Chem.* 280 (2005) 32230.
- [153] Y. Yao, A.V. Ferrer-Montiel, M. Montal, R.Y. Tsien, Activation of store-operated Ca²⁺ current in *Xenopus* oocytes requires SNAP-25 but not a diffusible messenger, *Cell* 98 (1999) 475.
- [154] A.S. Yoo, I. Cheng, S. Chung, T.Z. Grenfell, H. Lee, E. Pack-Chung, M. Handler, J. Shen, W. Xia, G. Tesco, A.J. Saunders, M.P. Frosch, R.E. Tanzi, T.-W. Kim, Presenilin-mediated modulation of capacitance calcium entry, *Neuron* 27 (2000) 561.
- [155] F. Zeng, S.Z. Xu, P.K. Jackson, D. McHugh, B. Kumar, S.J. Fountain, D. J. Beech, Human TRPC5 channel activated by a multiplicity of signals in a single cell, *J. Physiol.* 559 (2004) 739.
- [156] S.L. Zhang, A.V. Yeromin, X.H. Zhang, Y. Yu, O. Safrina, A. Penna, J. Roos, K.A. Stauderman, M.D. Cahalan, Genome-wide RNAi screen of Ca²⁺ influx identifies genes that regulate Ca²⁺ release-activated Ca²⁺ channel activity, *Proc. Natl. Acad. Sci. U. S. A.* 103 (2006) 9357.
- [157] S.L. Zhang, Y. Yu, J. Roos, J.A. Kozak, T.J. Deerinck, M.H. Ellisman, K. A. Stauderman, M.D. Cahalan, STIM1 is a Ca²⁺ sensor that activates CRAC channels and migrates from the Ca²⁺ store to the plasma membrane, *Nature* 437 (2005) 902.
- [158] Z. Zhang, Y. Tang, M.X. Zhu, Increased inwardly rectifying potassium currents in HEK-293 cells expressing murine transient receptor potential 4, *Biochem. J.* 354 (2001) 717.
- [159] X. Zhu, P.B. Chu, M. Peyton, L. Birnbaumer, Molecular cloning of a widely expressed human homologue for the *Drosophila* trp gene, *FEBS Lett.* 373 (1995) 193.
- [160] C. Zitt, A. Zobel, A.G. Obukhov, C. Harteneck, F. Kalkbrenner, A. Lückhoff, G. Schultz, Cloning and functional expression of a human Ca²⁺-permeable cation channel activated by calcium store depletion, *Neuron* 16 (1996) 1189.
- [161] F. Zufall, The TRPC2 ion channel and pheromone sensing in the accessory olfactory system, *Naunyn-Schmiedeberg Arch. Pharmacol.* 371 (2005) 245.
- [162] F. Zufall, K. Ukhanov, P. Lucas, T. Leinders-Zufall, Neurobiology of TRPC2: from gene to behavior, *Pflugers Arch.* 451 (2005) 61.
- [163] A. Zweifach, R.S. Lewis, Mitogen-regulated Ca²⁺ current of T lymphocytes is activated by depletion of intracellular Ca²⁺ stores, *Proc. Natl. Acad. Sci. U. S. A.* 90 (1993) 6295.