



Nursing Services Guidelines PACU/PAT/PEC

- Title:** **BLOOD GLUCOSE MONITORING PROTOCOL**
- Responsibility:** Anesthesia Department/Medical Staff
- Purpose of Guidelines:** To optimize glycemic control for surgical patients peri-operatively which will assist in minimizing post-operative infection and/or other complications.
- Procedure:**
- 1) Diabetic patients should be scheduled as a first case if possible.
 - 2) All diabetics will have an HbA1c ordered in the clinic, at the time of booking the case.
 - a) < 7% – Go
 - b) 7-9% – Go/No Go on a case by case basis as determined by anesthesia and surgery
 - c) >9% – **NO GO**
 - 3) All patients are to have a FSBS on day of service in pre-op.
 - a) BS<200mg/dl – Go
 - b) BS>200mg/dl but <300mg/dl – Go/No Go on a case by case basis as determined by anesthesia and surgery
 - c) BS>300mg/dl – **NO GO**
 - 4) Recommendations for insulin the day of service.
 - a) Patients should take their full basal insulin dose (long acting)
 - 5) Metformin
 - a) For surgeries including abdominal, thoracic and intracranial: Metformin, and combination drugs including Metformin, should be discontinued after the last usual dose the morning of the day before the scheduled surgery. Metformin should not be taken the morning of surgery.
 - b) All other surgeries: Metformin can be taken the night before surgery but should not be taken the morning of surgery.
 - 6) Patients with insulin pumps
 - a) Continue their basal rate during the perioperative period. If surgery interferes with the site or position, then turn off and convert to an insulin gtt.
 - 7) Target FSBS in the OR
 - a) For patients during the perioperative period a target FSBS of 140 – 180 for those on an insulin gtt.

Reviewed by: Andrew Casabianca, M.D. Chair, Department of Anesthesiology & Medical Director of Operative Services,
Toni Woodley, RN, Nurse Manager PACU, POH, & PAT

Approved: 3/18 (New)

Reviewed:

Revised:

Reviewed by Policy & Standard Committee, 3/18, 3/21