

# The Na/K-ATPase/Src complex and cardiotoxic steroid-activated protein kinase cascades

Zhichuan Li · Zijian Xie

Received: 31 October 2007 / Revised: 24 January 2008 / Accepted: 29 January 2008  
© Springer-Verlag 2008

**Abstract** The Na/K-ATPase was discovered by Skou in 1957. Since then, the efforts of numerous investigators have led to the following conclusions: (a) This enzyme is indeed the molecular machine for the ATP-dependent and -coupled transport of Na<sup>+</sup> and K<sup>+</sup> across the plasma membrane of a living cell in which such a process (sodium pump) is detected. (b) The Na/K-ATPase is also an important signal transducer that not only interacts and regulates protein kinases, but also functions as a scaffold, capable of bringing the effector and effectors together to form functional signalosomes. This minireview discusses the interaction between the Na/K-ATPase and Src to illustrate how a P-type ATPase can act as a receptor, converting a ligand-binding signal to the activation of protein kinase cascades and the generation of second messengers.

**Keywords** Na/K-ATPase · Tyrosine kinase · Protein phosphorylation · Receptor · Protein kinase

## Abbreviations

A domain	activation domain
ADPKD	autosomal dominant polycystic kidney disease
CD2	second cytosolic domain
CD3	third cytosolic domain
CTS	cardiotonic steroids
EGFR	epidermal growth factor receptor

ERK	extracellular signal-regulated protein kinase
FAK	focal adhesion kinase
FRET	fluorescence resonance energy transfer
GST	glutathione-S-transferase
GPCR	G protein-coupled receptor
IP3	inositol triphosphate
IP3R	IP3 receptor
IRBIT	IP <sub>3</sub> R binding protein released with inositol 1,4,5-trisphosphate
MBG	marinobufagenin
N domain	nucleotide binding domain
P domain	phosphorylation domain
PI3K	phosphatidylinositol 3-kinase
PKC	protein kinase C
PLC	phospholipase C
PTKs	protein tyrosine kinases
ROMK	renal outer medullary K channel
SERCA	sarcoplasmic reticulum Ca-ATPase

## Introduction

The P-type ATPases control ion fluxes across the cell membrane, whereas membrane receptors convert extracellular signals into the activation of protein kinase cascades as well as the generation of second messengers. When compared, these two groups of functionally divergent membrane proteins do share several common properties. First, many P-type ATPases, like the membrane receptors, interact with membrane and soluble proteins to form large protein complexes in a cell-specific manner. Second, they undergo ligand (substrate) binding-dependent conformational changes. These changes in protein conformation

Z. Li · Z. Xie (✉)  
Department of Physiology and Pharmacology,  
University of Toledo College of Medicine,  
Health Science Campus,  
Mail Stop 1008, 3000 Arlington Avenue,  
Toledo, OH 43614-2598, USA  
e-mail: zi-jian.xie@utoledo.edu

allow the movement of ions across the cell membrane. They could also alter the function of an interacting protein, and then generate an ion movement-independent cellular action. Consistently, recent studies have demonstrated that the Na/K-ATPase interacts with protein kinases, phosphatase, membrane transporters, and other cellular proteins and that these interactions make the Na/K-ATPase an important signal transducer. For instance, the interaction between the Na/K-ATPase and Src forms a receptor that can confer ligand-like effects of ouabain on protein tyrosine phosphorylation and subsequently on cell growth. This article will first review the background and several key findings that are relevant to the identification of this new cellular signaling mechanism, and then speculate on the potential molecular mechanisms that govern the formation of the Na/K-ATPase/Src receptor complex.

### The Na/K-ATPase

The Na/K-ATPase was discovered by Skou as an energy-transducing ion pump that converts the free energy of ATP into transmembrane ion gradients [95]. It belongs to a family of integral membrane proteins called P-type ATPases. The formation of a transient phosphorylated aspartate residue during the catalytic cycle is a hallmark of all P-type ATPase family members. In addition, they undergo a cation-dependent E1 to E2 conformational change that is fueled by ATP hydrolysis and coupled to the ion movement across the cell membrane (for detailed structure, function, and regulation of these ATPases, see reviews [48, 59, 71, 97]). The Na/K-ATPase consists of two non-covalently linked  $\alpha$  and  $\beta$  subunits [10, 48, 97]. FXFD proteins, a group of seven structurally similar polypeptides, are expressed in a tissue-specific manner and appear to act as a third subunit of the enzyme. The  $\alpha$  subunit (about 112 kDa) contains the ATP and other ligand-binding sites, and is considered as the “catalytic subunit”. The  $\beta$  subunit is essential for the assembly of a fully functional enzyme. Several  $\alpha$  and  $\beta$  subunits have been identified and functionally characterized [10, 48, 97]. The isoforms are expressed in a tissue-specific manner. The  $\alpha 1$  isoform is found in all cells. The  $\alpha 2$  and  $\alpha 3$  isoforms are expressed in skeletal muscle, neuronal tissue, and cardiac myocytes. The  $\alpha 4$  isoform is expressed in the testis and regulates sperm motility [10, 89]. Structural similarities have been found in the non-heavy metal subgroup of P-type ATPases that includes Na/K-ATPase and H/K-ATPase (group IIc) as well as sarcoplasmic–endoplasmic reticulum Ca-ATPase (SERCA, group IIa) [40, 44, 59, 71, 98]. The crystal structures of the SERCA1a at both E1 and E2 states have been resolved [38, 102, 103]. More recently, the structure of the Na/K-ATPase has also been revealed at the E2

conformation [77]. Based on these structures, the Na/K-ATPase  $\alpha 1$  subunit, like SERCA1a, contains three distinct functional domains [77, 98]. The A domain consists of the N-terminus and the second cytosolic domain (CD2) connected to transmembrane helices M2 and M3. The  $\alpha$  subunit also has the highly conserved phosphorylation (P) domain that is close to the membrane and a relatively isolated nucleotide binding (N) domain. It appears that the A domain rotates while the N domain closes up during the transport cycle, which opens and closes the A, N, and P domains in the E1 and the E2, respectively.

### Src, tyrosine phosphorylation and regulation of cell growth by cardiotonic steroids

*Cardiotonic steroids* A major difference between the Na/K-ATPase and other P-type ATPases is its ability to bind a class of chemicals called cardiotonic steroids (CTS). Although it remains to be firmly established, most biochemical studies have concluded that binding of CTS such as ouabain to the Na/K-ATPase (either phosphorylated or non-phosphorylated) will keep the enzyme at an E2-like close conformation [16, 48, 96]. This action of ouabain is known to inhibit the ion-pumping function of Na/K-ATPase. As discussed in the following section, it also activates the Na/K-ATPase-associated Src, resulting in stimulation of protein tyrosine phosphorylation. Chemically, CTS include plant-derived digitalis drugs such as digoxin and ouabain, and vertebrate-derived aglycones such as bufalin and marinobufagenin (MBG) [1, 91]. While digoxin has been used for more than 200 years to manage congestive heart failure and/or atria fibrillation, bufalin and MBG represent the major active components of a traditional Chinese medicine called Chan'su that is prescribed as a cardiotonic, diuretic, and anodyne agent. Although CTS have been considered only as drugs since their discovery, recent studies have identified both ouabain and MBG as endogenous steroids whose production and secretion are regulated by multiple physiological stimuli including adrenocorticotrophic hormone and angiotensin II [28, 36, 57, 61, 91]. Moreover, they are found to play an important role in the regulation of renal salt handling, vascular and cardiac contractions.

In addition to the aforementioned cardiac and vascular effects, CTS play an important role in the regulation of cell growth. In fact, the effects of ouabain on gene expression and the mitogen-induced differentiation and proliferation of lymphoblasts were first noted in the 1970s [24, 47, 85]. Recent studies have greatly expanded the diversity of CTS-affecting cells. Moreover, they showed that CTS activated multiple growth pathways and stimulated either differentiation/apoptosis or hypertrophic/proliferative growth in a cell-type-specific manner [9, 19, 26, 29–35, 41, 49, 56, 60,

62–65, 70, 73, 75, 79, 84, 87, 104]. Significantly, the growth regulatory effects of CTS could occur at nano- and sub-nanomolar concentrations that exhibited no inhibition of cellular Na/K-ATPase pumping activity [6, 53, 60, 64, 86, 90]. Finally, recent studies have demonstrated the potential physiological and pathological significance of CTS-mediated cell growth regulation [29, 50, 81]. To this end, it is important to note the following important studies: First, the effect of sub-nanomolar concentrations of ouabain on cell growth was first observed in cultured canine smooth muscle cells [6]. This observation was subsequently confirmed in several other types of cells including endothelial cells and cardiac fibroblasts [27, 64, 90]. Second, a recent study from Blanco's laboratory suggested that endogenous CTS might play an important role in the pathogenesis of autosomal dominant polycystic kidney disease (ADPKD) [81]. It is known that cyst formation and enlargement require proliferation of mural renal epithelial cells. Blanco and colleagues [81] found that a basal lateral application of sub-nanomolar concentrations of ouabain was sufficient to stimulate the proliferation of ADPKD cells, but not the normal human kidney cells, via the activation of an ERK pathway. Finally, recent *in vivo* studies have confirmed a role of CTS in the control of cardiac hypertrophy and fibrosis observed in uremic cardiomyopathy [27, 29, 50].

*Src and CTS-induced stimulation of protein tyrosine phosphorylation* The discovery of a class of proteins (e.g., v-Src and v-Abl) associated with the polyomavirus that phosphorylates tyrosine had revealed the importance of protein tyrosine phosphorylation in the control of cell growth [23]. Accordingly, many studies had been conducted to test whether ouabain and other CTS affected protein tyrosine phosphorylation [6, 22, 34, 55, 56, 58]. These studies demonstrated that sub-toxic concentrations of ouabain indeed stimulated tyrosine phosphorylation of multiple proteins in cardiac myocytes, A7r5 cells, HeLa, and LLC-PK1 cells. These ouabain effects were rapid and dose dependent. Moreover, the addition of a non-specific tyrosine kinase inhibitor such as genistein or herbimycin A was sufficient to block ouabain-induced tyrosine phosphorylation, and subsequently the growth effect of ouabain on cultured cells [6, 34]. More recently, these early observations were confirmed not only *in vitro* but also *in vivo* [29, 50]. It is interesting to note that studies have further revealed that increases in protein tyrosine phosphorylation are also important for CTS-induced cellular actions other than cell growth regulation [58, 100, 106, 113].


In principle, increases in protein tyrosine phosphorylation can occur through stimulation of protein tyrosine kinases or inhibition of tyrosine phosphatases or both. The Na/K-ATPase has no intrinsic tyrosine kinase activity. Thus, it is unlikely

that the Na/K-ATPase can serve as a receptor tyrosine kinase to convert the ouabain binding to increases in protein tyrosine phosphorylation. Although the  $\alpha$  subunit contains a consensus tyrosine phosphatase motif, preliminary studies have shown that it does not have significant tyrosine phosphatase activity (Tian and Xie, unpublished observation). There are many precedents where ligands stimulate protein tyrosine phosphorylation employing receptors that lack intrinsic kinase activity [20, 42, 74]. The examples of this mode of signal transduction include cytokine receptors [42] and G protein-coupled receptors (GPCRs) [74]. While the formation of “binary” cytokine receptors involves non-covalent association of JAK members with the receptor, the GPCRs employ Src family kinases. Thus, it is plausible that non-receptor tyrosine kinases such as Src could also be responsible for ouabain-induced protein tyrosine phosphorylation. Indeed, our initial studies revealed that ouabain increased translocation of cytosolic Src to a Triton-insoluble fraction and stimulated Src activity in cultured cardiac myocytes, A7r5, and LLC-PK1 cells [34]. The stimulation was apparently because of an increase in tyrosine phosphorylation of Src Y418, but not the dephosphorylation of Y529. Moreover, inhibition of Src by several inhibitors blocked ouabain-induced tyrosine phosphorylation of cellular proteins and abolished many ouabain-activated down-stream signal pathways including the activation of ERK. These findings were further confirmed in SYF mouse fibroblasts where Src family kinases were knocked out [35]. Functionally, activation of Src is not only important for CTS to regulate cell growth, but also other cellular activities [45, 52, 58, 67, 69, 82]. To this end, it is of interest to mention that the activated Src appears to be able to phosphorylate the Na/K-ATPase and then regulate the pump activity [11, 12, 29]. Historically, this mode of regulation had been suggested in the 1980s [80].

#### Identification of Na/K-ATPase/Src complex as a functional CTS receptor

*Structure and regulation of Src family kinases* Src family kinases are membrane-associated non-receptor tyrosine kinases. The prototype member of the Src family protein tyrosine kinases was first identified as v-Src, the molecule responsible for the cell-transforming ability of the Rous sarcoma virus [14]. So far, at least nine Src family members have been identified and can be divided into two groups: tyrosine kinases with a broad expression (Src, Fyn, Yes) and those mainly expressed in hematopoietic cells (Fgr, Lyn, Hck, Lck, Blk, Yrc, and Yrk) [99]. Structurally, each Src family kinase contains several distinct functional domains (Fig. 1a): (1) the N-terminal Src homology 4 (or SH4) domain, which is involved in targeting Src and other kinases to the membranes through myristoylation and/or palmitoyla-

**Fig. 1** **a** Schematic presentation of Na/K-ATPase and Src. The identified interacting domains are marked by *arrows*. **b** Modeling of Na/K-ATPase/Src interaction. Modeling of Na/K-ATPase E1- and E2-like structures was based on SERCA1a structure files 1SU4 and 1IWO as well as the newly published Na/K-ATPase structure, and then generated using SPDBView V3.7 program. The A domain (N-terminus and CD2) in Na/K-ATPase was labeled in *blue*, P domain in *green*, N domain in *black*. The SH2 domain of Src was labeled in *orange*, kinase domain in *light blue*. As indicated, we speculate that conformational change from the E1 to the E2 may release the kinase domain, thus activating Src



tion; (2) the SH3, which contains about 60 amino acids and binds the polyproline motifs; (3) the SH2 domain, which interacts with the phosphorylated tyrosine residues; (4) the SH2–Kinase linker region, which can form intramolecular interaction with SH3 and regulate kinase activity; (5) the conserved kinase domain (SH1); and (6) a short C-terminal regulatory tail.

The crystal structures of inactive Src and Hck as well as active Lck have been resolved, and together they provide vivid snapshots of the domain interactions that regulate Src activity [94, 111, 112]. Overall, the kinase domain of the Src family closely resembles that of protein kinase A [37, 54], which consists of an N-terminal and a larger C-terminal lobe linked by the activation loop. The formation of a salt bridge between E312 and K297 (human c-Src sequence) in the N lobe of the kinase domain is required for kinase activity. In addition, kinase activity is regulated by two important and highly conserved tyrosine residues (Y529 and Y418). Phosphorylation of Y529 leads to an intramolecular interaction between the SH2 domain and pY529, which facilitates binding of the SH3 domain to the linker region polyproline type II helix. This SH3-mediated interaction, in turn, inhibits the formation of the salt bridge between E312 and K297, and thus keeps Src in an inactive state. In accordance with the above model, Src can be activated when the interaction between the SH2 domain and pY529 is disrupted because of competitive binding of the

SH2 domain to the phosphotyrosine in other proteins such as EGFR. On the other hand, autophosphorylation of Y418 stabilizes the Src in an active state and then stimulates Src activity. Src is known to interact with a number of cellular proteins via various protein domains. The complexities of these interactions suggest that many cellular mechanisms exist for inhibiting/activating the kinase. Examples of Src activators include proteins interacting with the SH2 domain (e.g., platelet-derived growth factor receptor [3] and FAK [21]), SH3 domain (e.g., Nef [76, 105] and Sin [2]), and both SH2 and SH3 (e.g., p130<sup>CAS</sup> [15]).

*The interaction between the Na/K-ATPase and Src* The following evidence supports the notion that the Na/K-ATPase directly interacts with Src to form a functional receptor complex in live cells [58, 65, 68, 101, 107]: First, several laboratories, including ours, observed that the Na/K-ATPase and Src were co-enriched in caveolar fractions in different types of cells. Second, immunofluorescence imaging analysis showed co-localization of these two proteins in the plasma membrane. Third, both proteins could be co-immunoprecipitated by either anti- $\alpha 1$  or anti-Src antibodies. Fourth, fluorescence resonance energy transfer (FRET) analysis indicated that both proteins were in close proximity, providing further support of a direct interaction in live cells. Finally, in vitro GST pull-down assay demonstrated direct interactions between the  $\alpha 1$

subunit of Na/K-ATPase and Src. As depicted in Fig. 1a, the interaction between the  $\alpha 1$  and Src involves at least two contacting sites: one being the CD2 of  $\alpha 1$  and Src SH2, and the other consisting of the third cytosolic domain (CD3) of  $\alpha 1$  and the Src kinase domain. Based on the predicted structure of  $\alpha 1$ , the CD2 and CD3 constitute the A and N domains, respectively, and they are highly exposed in the E1 conformation (Fig. 1b). It is interesting to note that both domains are known to interact with structure, membrane, and soluble proteins. Examples of these interactions include ankyrin, inositol 1,4,5-trisphosphate (IP3) receptors (IP3Rs), phosphoinositide 3' kinase (PI3K), phospholipase C- $\gamma$  (PLC- $\gamma$ ), and cofilin [7, 43, 62, 113–116].

Functionally, the interaction between the Na/K-ATPase and Src keeps Src in an inactive state. When the molecular mechanism of this inhibition was probed, we found that the CD3/kinase domain interaction was responsible for this inhibition [101]. This is reminiscent of Wiscott–Aldrich syndrome protein-induced Src inhibition [92], but in contrast to the Src kinase domain interaction with G $\alpha$ s and G $\alpha$ i, which results in Src activation [72]. It is interesting to note that even though the RACK1/SH2 interaction inactivates Src [17, 18], the CD2/SH2 interaction appears to have no effect on the kinase activity. When the effect of ouabain on these domain interactions was examined, we observed that ouabain reduced the binding of the Src kinase domain, but not the SH2 or full-length Src, to the Na/K-ATPase. Thus, as depicted in Fig. 1b, we suggest that binding of ouabain to the Na/K-ATPase/Src receptor complex may free the Src kinase domain, resulting in activation of the Na/K-ATPase-associated Src kinase. While the Na/K-ATPase provides the ligand (CTS) binding site, the Na/K-ATPase-associated Src functions as a signal transducer, converting as well as amplifying the ligand-binding signal to increases in protein tyrosine phosphorylation. This notion is further supported by the following observations: First, ouabain reduced the FRET efficiency between the Na/K-ATPase and Src in live cells. Concomitantly, it stimulated cellular Src activity. Second, ouabain stimulated the tyrosine phosphorylation of multiple proteins that were either associated with or in close proximity to the Na/K-ATPase/Src receptor complex in a Src-dependent manner. It is of interest to point out that this mode of Src activation is unique and different from other known mechanisms of Src activation induced by either receptor tyrosine kinases or GPCRs. Specifically, it does not require dephosphorylation of pY529 or binding of the Src SH2 to a phosphorylated tyrosine [72, 76, 99]. Moreover, the interaction between the Na/K-ATPase and Src represents a novel regulation of cellular Src activity because of the involvement of both the SH2 and the kinase domains.

To seek additional evidence to support the notion that the Na/K-ATPase/Src complex serves as a CTS receptor, we

determined whether a graded-knockdown of cellular Na/K-ATPase could release the Na/K-ATPase-interacting Src, resulting in an increase in basal Src activity and a concomitant loss of ouabain-activated protein tyrosine phosphorylation [65]. Graded knockdown of cellular Na/K-ATPase was achieved by transfecting LLC-PK1 cells with a vector carrying  $\alpha 1$ -specific siRNA. When basal Src activity was measured, we found that knockdown of the Na/K-ATPase resulted in a significant increase in basal Src activity in a  $\alpha 1$  amount-dependent manner. Moreover, tyrosine phosphorylation of FAK, a Src effector, was also elevated. As expected, ouabain failed to stimulate Src and ERK1/2 in the knockdown cells. Furthermore, when the knockdown cells were rescued by rat  $\alpha 1$ , it not only restored basal Src/FAK activity, but also enabled ouabain to stimulate Src and ERK1/2 at a much higher concentration, consistent with the established differences in ouabain sensitivity between pig and rat  $\alpha 1$ . Finally, changes in cellular Src/FAK activity in the knockdown cells could also be rescued by expression of a pumping-null rat  $\alpha 1$  mutant. Both FRET analysis and co-immunoprecipitation assay showed that the pumping-null mutant was fully capable of interacting with Src. It is important to note that this mode of interaction appears to operate *in vivo* as well since Src activity in tissues harvested from  $\alpha 1^{+/-}$  mice were significantly increased (Li and Xie, unpublished data). Taken together, the data indicate that there is a pool of the Na/K-ATPase-interacting Src in live cells. The interaction between the Na/K-ATPase and Src not only regulates basal Src activity, but also serves as a function receptor for CTS to stimulate Src and subsequently protein tyrosine phosphorylation.

Despite progress in understanding the Na/K-ATPase/Src receptor complex, several important issues are worthy of further discussion. First, the Src family consists of at least nine members and many of them are expressed in a tissue-specific manner. Since these kinases are highly conserved, we suggest that the Na/K-ATPase may interact with other members of Src family kinases. We further speculate that these interactions may constitute the tissue-specific receptor Na/K-ATPase/Src family kinase complex that provides a certain degree of signaling specificity. Second, there are at least four  $\alpha$  isoforms. Because these isoforms are highly conserved, it is most likely that  $\alpha$  isoforms other than  $\alpha 1$  would interact with Src or Src family kinases to form a receptor complex. To this end, it is of interest to mention that Src also interacts with H/K-ATPase [101], potentially forming yet another group of receptor complexes. Moreover, recent studies have demonstrated that both Na/K-ATPase and H/K-ATPase are effectors of Src family kinases [46]. All together, the diversity of these interactions could produce a large population of receptor species and provide a tissue-specific response. Needless to say, these issues have to be further investigated.

*Signal amplification, conversion, and termination* Because CTS bind and consequently inhibit the pumping function of the Na/K-ATPase, they have been considered as specific pump inhibitors. Accordingly, the primary effects of CTS on various cellular functions have been ascribed to the inhibition of the Na/K-ATPase, and subsequent changes in intracellular ion concentration. In contrast, CTS would activate the Na/K-ATPase/Src receptor complex, thus making CTS true agonists of the receptor. Functionally, activation of this receptor complex by CTS results in signal amplification and conversion via the stimulation of protein kinase cascades and generation of second messengers. For example, it has been reported that binding of ouabain to this receptor complex was capable of recruiting and transactivating the EGFR [6, 34, 35, 58, 60]. This apparently involved Src-dependent phosphorylation of the EGFR at sites other than the major autophosphorylation site Y1173 [35]. The trans-activated EGFR, in turn, recruited adaptor protein Shc to the complex, resulting in activation of the Ras/Raf/MEK/ERK cascade [6, 35]. Similarly, Src activation also led to stimulation of PLC- $\gamma$  and subsequent activation of protein kinase C (PKC) isozymes and IP<sub>3</sub>-mediated Ca<sup>2+</sup> signaling [113]. Moreover, ouabain was found to stimulate PI3K, which works in concert with PKC, resulting in endocytosis of this receptor complex. This explains how cells terminate the ouabain-activated signal transduction [67]. In short, the identified Na/K-ATPase/Src complex appears to function as other receptor tyrosine kinases, capable of amplifying and converting signals as well as terminating the CTS-provoked signal transduction. Clearly, the identification of this new cellular signaling mechanism has provided new insight into the molecular action of CTS. It has also established a new target for developing novel agonists and antagonists of this receptor complex. To this end, recent studies have shown that some derivatives of ouabain such as rostafuroxin indeed antagonize the effect of ouabain on Src, but have no effect on Na/K-ATPase activity [29].

Although many laboratories have documented that changes in protein interaction are sufficient for CTS to provoke signal transduction, it is important to point out that these new findings do not exclude a role of the ion-pumping function of Na/K-ATPase in CTS-induced effects. Because the Na/K-ATPase is functionally coupled to many ion transporters such as the Na<sup>+</sup>/Ca<sup>2+</sup> exchanger, localized inhibition of the pumping function will modulate function of the Na<sup>+</sup>/Ca<sup>2+</sup> exchanger and increases intracellular Ca<sup>2+</sup>, especially in cells such as cardiac myocytes where the Na<sup>+</sup>/Ca<sup>2+</sup> exchanger is highly expressed. Thus, we believe that both processes could work in concert in the regulation of cellular function such as myocardial contractility. Indeed, recent studies have demonstrated that either knockout of Na<sup>+</sup>/Ca<sup>2+</sup> in cardiac myocytes or inhibition of the Src/PLC $\epsilon$

pathway is effective in blocking ouabain-induced inotropy [4, 51, 82, 88]. Moreover, since intracellular Na<sup>+</sup> regulates the conformation of the Na/K-ATPase (e.g., the E1 state), it is plausible that changes in intracellular Na<sup>+</sup> concentration could also regulate the formation of the Na/K-ATPase/Src complex, thus cellular Src activity. Clearly, these issues remain to be experimentally resolved.

It is important to note that many ion transporters and channels, like the Na/K-ATPase, are capable of forming signaling complexes and then serving as receptors or as signal integrators. Examples for such roles of ion-transporting proteins include, but are not limited to, the Na<sup>+</sup>/H<sup>+</sup> exchanger 1 [8], transient receptor potential channels [25, 108], the red blood cell anion exchanger band 3 [13], and IP<sub>3</sub>Rs [5, 83, 93]. For instance, IP<sub>3</sub>R interacts and keeps a protein called IP<sub>3</sub>R binding protein released with IP<sub>3</sub> (IRBIT) in an inactive state. Binding of IP<sub>3</sub> to the IP<sub>3</sub>R changes the conformation of the receptor, resulting in release of IRBIT from IP<sub>3</sub>R, which subsequently binds and stimulates the Na<sup>+</sup>/vldelimiterspaceHCO<sub>3</sub><sup>-</sup> co-transporter in renal epithelial cells [5, 93].

*Molecular mechanism of receptor formation* As depicted in Fig. 1b, modeling of interactions between the Na/K-ATPase and Src has led us to propose the following working hypotheses: First, we believe that both CD2/SH2 and N domain/kinase domain interactions could simultaneously occur because our in vitro studies showed that ouabain did not reduce binding of Src to the Na/K-ATPase [101]. Second, based on the crystal structures of Src, Na/K-ATPase, and SERCA, we believe that formation of the Na/K-ATPase/Src receptor complex would occur (or stabilize the Na/K-ATPase) in an E1-like conformation. We further suggest that the Src kinase domain would dissociate from the Na/K-ATPase in an E2-like conformation because of rotation of the A domain and the inward movement of the N domain (Fig. 1b). Specifically, we believe that this is how ouabain activates the Na/K-ATPase-associated Src. Third, we found that the CD2/SH2 interaction had a higher affinity than that of the N domain/kinase domain interaction (Li and Xie, unpublished data). Therefore, we contend that the CD2/SH2 interaction would facilitate the interaction between the N domain and the kinase domain, thus helping the formation of the Na/K-ATPase/Src receptor complex. Moreover, when ouabain turned the Na/K-ATPase into an E2-like conformation [16, 96], rotation of the A domain (i.e., CD2) would work in concert with the movement of the N domain, resulting in the release of the Src kinase domain and thus the activation of Src. Therefore, we propose that formation of a functional receptor Na/K-ATPase/Src complex requires both CD2/SH2 and N domain/kinase domain interactions. Needless to say, this proposal has to be experimentally tested.

## Conclusions and perspectives

Studies from many laboratories of the past few years have documented that Na/K-ATPase has a receptor function that confers a ligand-like effect of CTS on protein tyrosine phosphorylation in addition to the well-described ion-pumping function. Meanwhile, there is also increased appreciation of other membrane transporters in signal transduction. These studies certainly mark the beginning of a fascinating new field of investigation, evidenced by rapid growth of research concerning the role of these membrane proteins in signal transduction. Many important issues, especially those related to mechanism and physiological significance, remain to be resolved. For instance, does E1 and E2 conformational cycling regulate the interaction between the Na/K-ATPase and Src as predicted in Fig. 1b? Resolving this issue will certainly enhance our understanding of the molecular mechanism of CTS action. It may also reveal an intimate relationship between ion pumping and signal transduction. In principle, the Na/K-ATPase/Src complex could continuously generate signals through E1 and E2 cycling when it pumps. This mechanism, if proven to be true, would connect two basic cellular activities (pumping and signal transduction) together, which may be relevant to a well-documented cellular signaling mechanism of “pump–leak” coupling [39, 78, 109, 110]. It is known that apical and basolateral transport activities are tightly balanced, and changes in basolateral Na/K-ATPase pumping activity directly affect the apical cation conductance in renal epithelial cells. It was recently reported that activation of Src led to decreased exocytosis of renal outer medullary K channels (ROMK) to the apical membrane [66]. Thus, changes in the pumping activity (E1/E2 cycling) could affect Src activity, resulting in a feedback regulation of cellular leaks (e.g., ROMK-mediated K<sup>+</sup> efflux).

**Acknowledgments** We thank Ms. Martha Heck for editing the manuscript. This work was supported by NIH grants HL-36573 and HL-67963 awarded by the National Heart, Lung and Blood Institute and NIH grant GM-78565 awarded by the National Institute of General Medical Sciences.

## References

- Akera T, Brody TM (1976) Inotropic action of digitalis and ion transport. *Life Sci* 18:135–144
- Alexandropoulos K, Baltimore D (1996) Coordinate activation of c-Src by SH3- and SH2-binding sites on a novel p130Cas-related protein. *Sin. Genes Dev* 10:1341–1355
- Alonso G, Koegl M, Mazurenko N, Courtneidge SA (1995) Sequence requirements for binding of Src family tyrosine kinases to activated growth factor receptors. *J Biol Chem* 270:9840–9848
- Altamirano J, Li Y, DeSantiago J, Piacentino V 3rd, Houser SR, Bers DM (2006) The inotropic effect of cardioactive glycosides in ventricular myocytes requires Na<sup>+</sup>–Ca<sup>2+</sup> exchanger function. *J Physiol* 575:845–854
- Ando H, Mizutani A, Matsu-ura T, Mikoshiba K (2003) IRBIT, a novel inositol 1,4,5-trisphosphate (IP3) receptor-binding protein, is released from the IP3 receptor upon IP3 binding to the receptor. *J Biol Chem* 278:10602–10612
- Aydemir-Koksoy A, Abramowitz J, Allen JC (2001) Ouabain-induced signaling and vascular smooth muscle cell proliferation. *J Biol Chem* 276:46605–46611
- Barwe SP, Anilkumar G, Moon SY, Zheng Y, Whitelegge JP, Rajasekaran SA, Rajasekaran AK (2005) Novel role for Na,K-ATPase in phosphatidylinositol 3-kinase signaling and suppression of cell motility. *Mol Biol Cell* 16:1082–1094
- Baumgartner M, Patel H, Barber DL (2004) Na(+)/H(+) exchanger NHE1 as plasma membrane scaffold in the assembly of signaling complexes. *Am J Physiol Cell Physiol* 287:C844–850
- Bielawski K, Winnicka K, Bielawska A (2006) Inhibition of DNA topoisomerases I and II, and growth inhibition of breast cancer MCF-7 cells by ouabain, digoxin and proscillaridin A. *Biol Pharm Bull* 29:1493–1497
- Blanco G, Mercer RW (1998) Isozymes of the Na–K-ATPase: heterogeneity in structure, diversity in function. *Am J Physiol* 275:F633–650
- Bozulic LD, Dean WL, Delamere NA (2004) The influence of Lyn kinase on Na,K-ATPase in porcine lens epithelium. *Am J Physiol Cell Physiol* 286:C90–96
- Bozulic LD, Dean WL, Delamere NA (2005) The influence of SRC-family tyrosine kinases on Na,K-ATPase activity in lens epithelium. *Invest Ophthalmol Vis Sci* 46:618–622
- Bruce LJ, Beckmann R, Ribeiro ML, Peters LL, Chasis JA, Delaunay J, Mohandas N, Anstee DJ, Tanner MJ (2003) A band 3-based macrocomplex of integral and peripheral proteins in the RBC membrane. *Blood* 101:4180–4188
- Brugge JS, Erikson RL (1977) Identification of a transformation-specific antigen induced by an avian sarcoma virus. *Nature* 269:346–348
- Burnham MR, Bruce-Staskal PJ, Harte MT, Weidow CL, Ma A, Weed SA, Bouton AH (2000) Regulation of c-SRC activity and function by the adapter protein CAS. *Mol Cell Biol* 20:5865–5878
- Carilli CT, Farley RA, Perlman DM, Cantley LC (1982) The active site structure of Na<sup>+</sup>- and K<sup>+</sup>-stimulated ATPase. Location of a specific fluorescein isothiocyanate reactive site. *J Biol Chem* 257:5601–5606
- Chang BY, Chiang M, Cartwright CA (2001) The interaction of Src and RACK1 is enhanced by activation of protein kinase C and tyrosine phosphorylation of RACK1. *J Biol Chem* 276:20346–20356
- Chang BY, Conroy KB, Machleder EM, Cartwright CA (1998) RACK1, a receptor for activated C kinase and a homolog of the beta subunit of G proteins, inhibits activity of src tyrosine kinases and growth of NIH 3T3 cells. *Mol Cell Biol* 18:3245–3256
- Chueh SC, Guh JH, Chen J, Lai MK, Teng CM (2001) Dual effects of ouabain on the regulation of proliferation and apoptosis in human prostatic smooth muscle cells. *J Urol* 166:347–353
- Clements JL, Koretzky GA (1999) Recent developments in lymphocyte activation: linking kinases to downstream signaling events. *J Clin Invest* 103:925–929
- Cobb BS, Schaller MD, Leu TH, Parsons JT (1994) Stable association of pp60src and pp59fyn with the focal adhesion-associated protein tyrosine kinase, pp125FAK. *Mol Cell Biol* 14:147–155
- Contreras RG, Shoshani L, Flores-Maldonado C, Lazaro A, Cerejido M (1999) Relationship between Na(+),K(+)-ATPase and cell attachment. *J Cell Sci* 112(Pt 23):4223–4232

23. Courtneidge SA, Smith AE (1983) Polyoma virus transforming protein associates with the product of the c-src cellular gene. *Nature* 303:435–439
24. Cuff JM, Lichtman A (1975) The early effects of ouabain on potassium metabolism and rate of proliferation of mouse lymphoblasts. *J Cell Physiol* 85:209–215
25. Delmas P (2005) Polycystins: polymodal receptor/ion-channel cellular sensors. *Pflugers Arch* 451:264–276
26. Dmitrieva RI, Doris PA (2003) Ouabain is a potent promoter of growth and activator of ERK1/2 in ouabain-resistant rat renal epithelial cells. *J Biol Chem* 278:28160–28166
27. Elkareh J, Kennedy DJ, Yashaswi B, Vetteth S, Shidyak A, Kim EG, Smaili S, Periyasamy SM, Harii IM, Fedorova L, Liu J, Wu L, Kahaleh MB, Xie Z, Malhotra D, Fedorova OV, Kashkin VA, Bagrov AY, Shapiro JI (2007) Marinobufagenin stimulates fibroblast collagen production and causes fibrosis in experimental uremic cardiomyopathy. *Hypertension* 49:215–224
28. Fedorova OV, Doris PA, Bagrov AY (1998) Endogenous marinobufagenin-like factor in acute plasma volume expansion. *Clin Exp Hypertens* 20:581–591
29. Ferrandi M, Molinari I, Barassi P, Minotti E, Bianchi G, Ferrari P (2004) Organ hypertrophic signaling within caveolae membrane subdomains triggered by ouabain and antagonized by PST 2238. *J Biol Chem* 279:33306–33314
30. Feschenko MS, Wetzel RK, Sweadner KJ (1997) Phosphorylation of Na,K-ATPase by protein kinases. Sites, susceptibility, and consequences. *Ann N Y Acad Sci* 834:479–488
31. Golden WC, Martin LJ (2006) Low-dose ouabain protects against excitotoxic apoptosis and up-regulates nuclear Bcl-2 in vivo. *Neuroscience* 137:133–144
32. Golomb E, Hill MR, Brown RG, Keiser HR (1994) Ouabain enhances the mitogenic effect of serum in vascular smooth muscle cells. *Am J Hypertens* 7:69–74
33. Griffiths NM, Ogden PH, Cormack R, Lamb JF (1991) Discrepancy between the short and long term effects of ouabain on the sodium pumps of human cells grown in culture. *Br J Pharmacol* 104:419–427
34. Haas M, Askari A, Xie Z (2000) Involvement of Src and epidermal growth factor receptor in the signal-transducing function of Na<sup>+</sup>/K<sup>+</sup>-ATPase. *J Biol Chem* 275:27832–27837
35. Haas M, Wang H, Tian J, Xie Z (2002) Src-mediated inter-receptor cross-talk between the Na<sup>+</sup>/K<sup>+</sup>-ATPase and the epidermal growth factor receptor relays the signal from ouabain to mitogen-activated protein kinases. *J Biol Chem* 277:18694–18702
36. Hamlyn JM, Blaustein MP, Bova S, DuCharme DW, Harris DW, Mandel F, Mathews WR, Ludens JH (1991) Identification and characterization of a ouabain-like compound from human plasma. *Proc Natl Acad Sci U S A* 88:6259–6263
37. Hanks SK, Quinn AM, Hunter T (1988) The protein kinase family: conserved features and deduced phylogeny of the catalytic domains. *Science* 241:42–52
38. Hilge M, Siegal G, Vuister GW, Guntert P, Gloor SM, Abrahams JP (2003) ATP-induced conformational changes of the nucleotide-binding domain of Na,K-ATPase. *Nat Struct Biol* 10:468–474
39. Hoffmann EK (2001) The pump and leak steady-state concept with a variety of regulated leak pathways. *J Membr Biol* 184:321–330
40. Horisberger JD (2004) Recent insights into the structure and mechanism of the sodium pump. *Physiology (Bethesda)* 19:377–387
41. Huang L, Li H, Xie Z (1997) Ouabain-induced hypertrophy in cultured cardiac myocytes is accompanied by changes in expression of several late response genes. *J Mol Cell Cardiol* 29:429–437
42. Ihle JN, Kerr IM (1995) Jaks and Stats in signaling by the cytokine receptor superfamily. *Trends Genet* 11:69–74
43. Jordan C, Puschel B, Koob R, Drenckhahn D (1995) Identification of a binding motif for ankyrin on the alpha-subunit of Na<sup>+</sup>, K<sup>+</sup>-ATPase. *J Biol Chem* 270:29971–29975
44. Jorgensen PL, Hakansson KO, Karlsh SJ (2003) Structure and mechanism of Na,K-ATPase: functional sites and their interactions. *Annu Rev Physiol* 65:817–849
45. Jung J, Kim M, Choi S, Kim MJ, Suh JK, Choi EC, Lee K (2006) Molecular mechanism of cofilin dephosphorylation by ouabain. *Cell Signal* 18:2033–2040
46. Kanagawa M, Watanabe S, Kaya S, Togawa K, Imagawa T, Shimada A, Kikuchi K, Taniguchi K (2000) Membrane enzyme systems responsible for the Ca(2+)-dependent phosphorylation of Ser(27), the independent phosphorylation of Tyr(10) and Tyr(7), and the dephosphorylation of these phosphorylated residues in the alpha-chain of H/K-ATPase. *J Biochem (Tokyo)* 127:821–828
47. Kaplan JG (1978) Membrane cation transport and the control of proliferation of mammalian cells. *Annu Rev Physiol* 40:19–41
48. Kaplan JH (2002) Biochemistry of Na,K-ATPase. *Annu Rev Biochem* 71:511–535
49. Kawazoe N, Watabe M, Masuda Y, Nakajo S, Nakaya K (1999) Tiam1 is involved in the regulation of bufalin-induced apoptosis in human leukemia cells. *Oncogene* 18:2413–2421
50. Kennedy DJ, Vetteth S, Periyasamy SM, Kanj M, Fedorova L, Khouri S, Kahaleh MB, Xie Z, Malhotra D, Kolodkin NI, Lakatta EG, Fedorova OV, Bagrov AY, Shapiro JI (2006) Central role for the cardiotonic steroid marinobufagenin in the pathogenesis of experimental uremic cardiomyopathy. *Hypertension* 47:488–495
51. Kennedy DJ, Vetteth S, Xie M, Periyasamy SM, Xie Z, Han C, Basrur V, Mutgi K, Fedorov V, Malhotra D, Shapiro JI (2006) Ouabain decreases sarco(endo)plasmic reticulum calcium ATPase activity in rat hearts by a process involving protein oxidation. *Am J Physiol Heart Circ Physiol* 291:H3003–3011
52. Khundmiri SJ, Amin V, Henson J, Lewis J, Ameen M, Rane MJ, Delamere NA (2007) Ouabain stimulates protein kinase B (Akt) phosphorylation in opossum kidney proximal tubule cells through an ERK-dependent pathway. *Am J Physiol Cell Physiol* 293:C1171–1180
53. Khundmiri SJ, Metzler MA, Ameen M, Amin V, Rane MJ, Delamere NA (2006) Ouabain induces cell proliferation through calcium-dependent phosphorylation of Akt (protein kinase B) in opossum kidney proximal tubule cells. *Am J Physiol Cell Physiol* 291:C1247–1257
54. Knighton DR, Zheng JH, Ten Eyck LF, Ashford VA, Xuong NH, Taylor SS, Sowadski JM (1991) Crystal structure of the catalytic subunit of cyclic adenosine monophosphate-dependent protein kinase. *Science* 253:407–414
55. Kometiani P, Li J, Gnudi L, Kahn BB, Askari A, Xie Z (1998) Multiple signal transduction pathways link Na<sup>+</sup>/K<sup>+</sup>-ATPase to growth-related genes in cardiac myocytes. The roles of Ras and mitogen-activated protein kinases. *J Biol Chem* 273:15249–15256
56. Kometiani P, Liu L, Askari A (2005) Digitalis-induced signaling by Na<sup>+</sup>/K<sup>+</sup>-ATPase in human breast cancer cells. *Mol Pharmacol* 67:929–936
57. Komiya Y, Dong XH, Nishimura N, Masaki H, Yoshika M, Masuda M, Takahashi H (2005) A novel endogenous digitalis, telocinobufagin, exhibits elevated plasma levels in patients with terminal renal failure. *Clin Biochem* 38:36–45
58. Kotova O, Al-Khalili L, Talia S, Hooke C, Fedorova OV, Bagrov AY, Chibalin AV (2006) Cardiotonic steroids stimulate glycogen synthesis in human skeletal muscle cells via a Src- and ERK1/2-dependent mechanism. *J Biol Chem* 281:20085–20094
59. Kuhlbrandt W (2004) Biology, structure and mechanism of P-type ATPases. *Nat Rev Mol Cell Biol* 5:282–295
60. Kulikov A, Eva A, Kirch U, Boldyrev A, Scheiner-Bobis G (2007) Ouabain activates signaling pathways associated with cell

- death in human neuroblastoma. *Biochim Biophys Acta* 1768:1691–1702
61. Laredo J, Shah JR, Lu ZR, Hamilton BP, Hamlyn JM (1997) Angiotensin II stimulates secretion of endogenous ouabain from bovine adrenocortical cells via angiotensin type 2 receptors. *Hypertension* 29:401–407
  62. Lee K, Jung J, Kim M, Guidotti G (2001) Interaction of the alpha subunit of Na,K-ATPase with cofilin. *Biochem J* 353:377–385
  63. Li J, Zelenin S, Aperia A, Aizman O (2006) Low doses of ouabain protect from serum deprivation-triggered apoptosis and stimulate kidney cell proliferation via activation of NF- $\kappa$ B. *J Am Soc Nephrol* 17:1848–1857
  64. Li M, Wang Q, Guan L (2007) Effects of ouabain on proliferation, intracellular free calcium and c-myc mRNA expression in vascular smooth muscle cells. *J Comp Physiol [B]* 177:589–595
  65. Liang M, Cai T, Tian J, Qu W, Xie ZJ (2006) Functional characterization of Src-interacting Na/K-ATPase using RNA interference assay. *J Biol Chem* 281:19709–19719
  66. Lin DH, Sterling H, Yang B, Hebert SC, Giebisch G, Wang WH (2004) Protein tyrosine kinase is expressed and regulates ROMK1 location in the cortical collecting duct. *Am J Physiol Renal Physiol* 286:F881–892
  67. Liu J, Kesiry R, Periyasamy SM, Malhotra D, Xie Z, Shapiro JJ (2004) Ouabain induces endocytosis of plasmalemmal Na/K-ATPase in LLC-PK1 cells by a clathrin-dependent mechanism. *Kidney Int* 66:227–241
  68. Liu L, Mohammadi K, Aynafshar B, Wang H, Li D, Liu J, Ivanov AV, Xie Z, Askari A (2003) Role of caveolae in signal-transducing function of cardiac Na<sup>+</sup>/K<sup>+</sup>-ATPase. *Am J Physiol Cell Physiol* 284:C1550–1560
  69. Liu L, Zhao X, Pierre SV, Askari A (2007) Association of PI3K/Akt signaling pathway with digitalis-induced hypertrophy of cardiac myocytes. *Am J Physiol Cell Physiol* 293:C1489–C1497
  70. Lopez-Lazaro M, Pastor N, Azrak SS, Ayuso MJ, Austin CA, Cortes F (2005) Digitoxin inhibits the growth of cancer cell lines at concentrations commonly found in cardiac patients. *J Nat Prod* 68:1642–1645
  71. Lutsenko S, Kaplan JH (1995) Organization of P-type ATPases: significance of structural diversity. *Biochemistry* 34:15607–15613
  72. Ma YC, Huang J, Ali S, Lowry W, Huang XY (2000) Src tyrosine kinase is a novel direct effector of G proteins. *Cell* 102:635–646
  73. McConkey DJ, Lin Y, Nutt LK, Ozel HZ, Newman RA (2000) Cardiac glycosides stimulate Ca<sup>2+</sup> increases and apoptosis in androgen-independent, metastatic human prostate adenocarcinoma cells. *Cancer Res* 60:3807–3812
  74. McGarrigle D, Huang XY (2007) GPCRs signaling directly through Src-family kinases. *Sci STKE* 2007:pe35
  75. Mijatovic T, Roland I, Van Quaquebeke E, Nilsson B, Mathieu A, Van Vynckt F, Darro F, Blanco G, Facchini V, Kiss R (2007) The alpha1 subunit of the sodium pump could represent a novel target to combat non-small cell lung cancers. *J Pathol* 212:170–179
  76. Moarefi I, LaFevre-Bernt M, Sicheri F, Huse M, Lee CH, Kuriyan J, Miller WT (1997) Activation of the Src-family tyrosine kinase Hck by SH3 domain displacement. *Nature* 385:650–653
  77. Morth JP, Pedersen BP, Toustrup-Jensen MS, Sorensen TL, Petersen J, Andersen JP, Vilsen B, Nissen P (2007) Crystal structure of the sodium-potassium pump. *Nature* 450:1043–1049
  78. Muto S, Asano Y, Seldin D, Giebisch G (1999) Basolateral Na<sup>+</sup> pump modulates apical Na<sup>+</sup> and K<sup>+</sup> conductances in rabbit cortical collecting ducts. *Am J Physiol* 276:F143–158
  79. Nakagawa Y, Rivera V, Larner AC (1992) A role for the Na/K-ATPase in the control of human *c-fos* and *c-jun* transcription. *J Biol Chem* 267:8785–8788
  80. Nakamura S, Braun S, Racker E (1987) A tyrosine-specific protein kinase from Ehrlich ascites tumor cells. *Arch Biochem Biophys* 252:538–548
  81. Nguyen AN, Wallace DP, Blanco G (2007) Ouabain binds with high affinity to the Na,K-ATPase in human polycystic kidney cells and induces extracellular signal-regulated kinase activation and cell proliferation. *J Am Soc Nephrol* 18:46–57
  82. Pasdois P, Quinlan CL, Rissa A, Tariosse L, Vinassa B, Costa AD, Pierre SV, Dos Santos P, Garlid KD (2007) Ouabain protects rat hearts against ischemia-reperfusion injury via pathway involving src kinase, mitoKATP, and ROS. *Am J Physiol Heart Circ Physiol* 292:H1470–1478
  83. Patterson RL, Boehning D, Snyder SH (2004) Inositol 1,4,5-trisphosphate receptors as signal integrators. *Annu Rev Biochem* 73:437–465
  84. Peng M, Huang L, Xie Z, Huang WH, Askari A (1996) Partial inhibition of Na<sup>+</sup>/K<sup>+</sup>-ATPase by ouabain induces the Ca<sup>2+</sup>-dependent expressions of early-response genes in cardiac myocytes. *J Biol Chem* 271:10372–10378
  85. Pollack LR, Tate EH, Cook JS (1981) Na<sup>+</sup>, K<sup>+</sup>-ATPase in HeLa cells after prolonged growth in low K<sup>+</sup> or ouabain. *J Cell Physiol* 106:85–97
  86. Qiu J, Gao HQ, Zhou RH, Liang Y, Zhang XH, Wang XP, You BA, Cheng M (2007) Proteomics analysis of the proliferative effect of low-dose ouabain on human endothelial cells. *Biol Pharm Bull* 30:247–253
  87. Ramirez-Ortega M, Maldonado-Lagunas V, Melendez-Zajgla J, Carrillo-Hernandez JF, Pastelin-Hernandez G, Picazo-Picazo O, Ceballos-Reyes G (2006) Proliferation and apoptosis of HeLa cells induced by in vitro stimulation with digitalis. *Eur J Pharmacol* 534:71–76
  88. Reuter H, Henderson SA, Han T, Ross RS, Goldhaber JL, Philipson KD (2002) The Na<sup>+</sup>-Ca<sup>2+</sup> exchanger is essential for the action of cardiac glycosides. *Circ Res* 90:305–308
  89. Sanchez G, Nguyen AN, Timmerberg B, Tash JS, Blanco G (2006) The Na,K-ATPase alpha4 isoform from humans has distinct enzymatic properties and is important for sperm motility. *Mol Hum Reprod* 12:565–576
  90. Saunders R, Scheiner-Bobis G (2004) Ouabain stimulates endothelin release and expression in human endothelial cells without inhibiting the sodium pump. *Eur J Biochem* 271:1054–1062
  91. Schoner W, Scheiner-Bobis G (2007) Endogenous and exogenous cardiac glycosides: their roles in hypertension, salt metabolism, and cell growth. *Am J Physiol Cell Physiol* 293:C509–C536
  92. Schulte RJ, Sefton BM (2003) Inhibition of the activity of SRC and Abl tyrosine protein kinases by the binding of the Wiskott–Aldrich syndrome protein. *Biochemistry* 42:9424–9430
  93. Shirakabe K, Priori G, Yamada H, Ando H, Horita S, Fujita T, Fujimoto I, Mizutani A, Seki G, Mikoshiba K (2006) IRBIT, an inositol 1,4,5-trisphosphate receptor-binding protein, specifically binds to and activates pancreas-type Na<sup>+</sup>/HCO<sub>3</sub><sup>-</sup> cotransporter 1 (pNBC1). *Proc Natl Acad Sci U S A* 103:9542–9547
  94. Sicheri F, Moarefi I, Kuriyan J (1997) Crystal structure of the Src family tyrosine kinase Hck. *Nature* 385:602–609
  95. Skou JC (1957) The influence of some cations on an adenosine triphosphatase from peripheral nerves. *Biochim Biophys Acta* 23:394–401
  96. Suzuki K, Taniguchi K, Iida S (1985) Ouabain binding and the conformational change in Na<sup>+</sup>, K<sup>+</sup>-ATPase. *Nippon Yakurigaku Zasshi* 86:181–188
  97. Sweadner KJ (1989) Isozymes of the Na<sup>+</sup>/K<sup>+</sup>-ATPase. *Biochim Biophys Acta* 988:185–220
  98. Sweadner KJ, Donnet C (2001) Structural similarities of Na,K-ATPase and SERCA, the Ca(2+)-ATPase of the sarcoplasmic reticulum. *Biochem J* 356:685–704

99. Thomas SM, Brugge JS (1997) Cellular functions regulated by Src family kinases. *Annu Rev Cell Dev Biol* 13:513–609
100. Thundathil JC, Anzar M, Buhr MM (2006) Na<sup>+</sup>/K<sup>+</sup> ATPase as a signaling molecule during bovine sperm capacitation. *Biol Reprod* 75:308–317
101. Tian J, Cai T, Yuan Z, Wang H, Liu L, Haas M, Maksimova E, Huang XY, Xie ZJ (2006) Binding of Src to Na<sup>+</sup>/K<sup>+</sup>-ATPase forms a functional signaling complex. *Mol Biol Cell* 17:317–326
102. Toyoshima C, Nakasako M, Nomura H, Ogawa H (2000) Crystal structure of the calcium pump of sarcoplasmic reticulum at 2.6 Å resolution. *Nature* 405:647–655
103. Toyoshima C, Nomura H (2002) Structural changes in the calcium pump accompanying the dissociation of calcium. *Nature* 418:605–611
104. Trevisi L, Visentin B, Cusinato F, Pighin I, Luciani S (2004) Antiapoptotic effect of ouabain on human umbilical vein endothelial cells. *Biochem Biophys Res Commun* 321:716–721
105. Tribble RP, Emert-Sedlak L, Smithgall TE (2006) HIV-1 Nef selectively activates Src family kinases Hck, Lyn, and c-Src through direct SH3 domain interaction. *J Biol Chem* 281:27029–27038
106. Valente RC, Capella LS, Monteiro RQ, Rumjanek VM, Lopes AG, Capella MA (2003) Mechanisms of ouabain toxicity. *FASEB J* 17:1700–1702
107. Wang H, Haas M, Liang M, Cai T, Tian J, Li S, Xie Z (2004) Ouabain assembles signaling cascades through the caveolar Na<sup>+</sup>/K<sup>+</sup>-ATPase. *J Biol Chem* 279:17250–17259
108. Wang T, Jiao Y, Montell C (2005) Dissecting independent channel and scaffolding roles of the Drosophila transient receptor potential channel. *J Cell Biol* 171:685–694
109. Wang WH, Geibel J, Giebisch G (1993) Mechanism of apical K<sup>+</sup> channel modulation in principal renal tubule cells. Effect of inhibition of basolateral Na(+)-K(+)-ATPase. *J Gen Physiol* 101:673–694
110. Welling PA (1995) Cross-talk and the role of KATP channels in the proximal tubule. *Kidney Int* 48:1017–1023
111. Xu W, Harrison SC, Eck MJ (1997) Three-dimensional structure of the tyrosine kinase c-Src. *Nature* 385:595–602
112. Yamaguchi H, Hendrickson WA (1996) Structural basis for activation of human lymphocyte kinase Lck upon tyrosine phosphorylation. *Nature* 384:484–489
113. Yuan Z, Cai T, Tian J, Ivanov AV, Giovannucci DR, Xie Z (2005) Na/K-ATPase tethers phospholipase C and IP3 receptor into a calcium-regulatory complex. *Mol Biol Cell* 16:4034–4045
114. Yudowski GA, Efendiev R, Pedemonte CH, Katz AI, Berggren PO, Bertorello AM (2000) Phosphoinositide-3 kinase binds to a proline-rich motif in the Na<sup>+</sup>, K<sup>+</sup>-ATPase alpha subunit and regulates its trafficking. *Proc Natl Acad Sci U S A* 97:6556–6561
115. Zhang S, Malmersjo S, Li J, Ando H, Aizman O, Uhlen P, Mikoshiba K, Aperia A (2006) Distinct role of the N-terminal tail of the Na,K-ATPase catalytic subunit as a signal transducer. *J Biol Chem* 281:21954–21962
116. Zhang Z, Devarajan P, Dorfman AL, Morrow JS (1998) Structure of the ankyrin-binding domain of alpha-Na,K-ATPase. *J Biol Chem* 273:18681–18684