

Ph.D. Qualifying Examination  
Spring 2003

**Instructions:**

1. If you think that there is a mistake ask the proctor. If the proctor's explanation is not satisfactory, interpret the problem as you see fit, but not in such a way that the answer is trivial.
2. From each part solve 3 of 4 problems.
3. If you solve more than three problems from a part indicate the problems that you wish to have graded.

**Part A: ODE Questions**

1. Suppose that  $g(v)$  is a continuously differentiable function from  $R^n$  to  $R^n$  satisfying  $\|Dg(v)\| \leq 2M$  and  $\|g(v)\| < M\|v\|$  for some positive constant  $M$ . Consider the initial value problem

$$\dot{v} = h + g(v); \quad v(0) = 0$$

where  $h$  is a constant vector in  $R^n$ . Show that the solution for  $t > 0$ ,  $v(t)$  satisfies

$$\|v(t) - ht\| \leq \frac{\|h\|}{4M^2}(e^{2Mt} - (2Mt + 1)).$$

2. Find the fundamental solution of  $X(t)$  with  $X(0) = I$  to the system

$$\frac{d}{dt} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} -x_2 + x_3 \\ x_1 + x_4 \\ -x_4 \\ x_3 \end{pmatrix}.$$

3. Consider the system  $\dot{x} = h(t)Ax$  where  $x : R \rightarrow R^n$ ,  $A$  is a constant  $n \times n$  matrix, and  $h(t)$  is strictly positive and continuous. Show that 0 is asymptotically stable if all the eigenvalues of  $A$  have negative real parts and  $\int_0^\infty h(t)dt$  diverges. By example show that the stability may not hold if the integral converges.

4. Let  $L$  be a periodic solution of a Lipschitz continuous planar autonomous system  $\dot{x} = f(x)$  with flow  $\phi_t : R^2 \rightarrow R^2$ . Recall that an  $\epsilon$ -neighborhood  $N_\epsilon$  of  $L$  is small if  $N_\epsilon$  contains no singular points and for any  $q_1, q_2 \in N_\epsilon$  with  $|q_1 - q_2| < 2\epsilon$ ,

$$\frac{\langle f(q_1), f(q_2) \rangle}{\|f(q_1)\| \|f(q_2)\|} > \frac{1}{\sqrt{2}}.$$

Suppose that  $L^1 \subset N_\epsilon$  is a second periodic solution and that for some  $p \in L$  and transverse segment  $\overline{n^1 p n}$  to  $L$  there is a  $p^1 \in L^1$  with  $p^1 \in \overline{n^1 p n}$ . Let  $T$  be the smallest parameter value so that  $\phi_T(p^1) \in \overline{n^1 p n}$ . Show that  $T$  is the period of  $L^1$ .

### Part B: PDE Questions

1. Consider the quasi-linear system for an unknown function  $u : R^n \times R \rightarrow R$  given by

$$\frac{\partial u}{\partial t} + \vec{a}(\vec{x}, t) \cdot \frac{\partial u}{\partial \vec{x}} = b(\vec{x}, t, u)$$

where  $t \in R$  and  $\vec{x} \in R^n$ . Show that if  $\vec{a}, b$  are bounded and Lipschitz continuous, then the Cauchy problem with Cauchy data defined on the hyperplane  $H = \{(\vec{x}, t) \mid t = 0\}$  has global solutions.

2. Determine the canonical form and the general solution to the second order equation

$$4x^2 u_{xx} + 4xy u_{xy} - 8y^2 u_{yy} + 4xu_x - 8yu_y = 0.$$

3. Suppose that on a bounded domain  $\Omega$  a function  $u \in C^2(\Omega) \cap C^0(\bar{\Omega})$  satisfies  $\Delta u + \lambda u^3 = 0$  in  $\Omega$  with  $\lambda > 0$  and  $u = 0$  on  $\partial\Omega$ . Suppose that  $u$  is non-negative in  $\Omega$ . Show that  $u$  is strictly positive in  $\Omega$ .

4. Suppose  $w$  is a harmonic function on  $R^n$  and satisfies

$$\int_{R^n} w^2(x) dx < \infty.$$

Prove that  $w \equiv 0$ .