

Nursing Service Guidelines General

Title: INDWELLING FOLEY CATHETER CARE

Responsibility: Registered Nurse (RN), or Licensed Practical Nurse (LPN), or Nursing Assistant (NA), or

Patient Care Technician (PCT)

Purpose:

To increase clinical knowledge on care of indwelling foley catheters and prevent catheter -

associated urinary tract infections (CAUTI).

Procedure:

1. Wash hands before and after touching any part of the catheter system and before and after applying gloves.

- 2. Check for proper drainage whenever coming in contact with patient, at least every four hours.
- 3. Keep the drainage bag below bladder level to prevent reflux of urine into the bladder.
- 4. Empty drainage bag at a minimum of every shift.
- 5. Keep the drainage tubing/catheter junction closed. If the catheter/tubing junction is broken, replace the catheter and collection system. Never clamp the drainage tubing except under doctor's order.
- 6. Perineal care needs to be done at the start of the day shift (0700-1900) and the start of the night shift (1900-0700).
- 7. Perineal care needs to be done immediately after bowel movements to ensure the perineal area is always clean.
- 8. The catheter site, and urine need to be inspected for potential problems, and problems need to be reported to the physician.
- 9. Foley catheter assessment needs to be completed each shift and a maintain order needs completed with patient meets the Nurse Driven Protocol criteria for use. Adequate fluid intake helps flush the urinary system and reduces the risk of infection.

Procedure

Point of Emphasis

- 1. Perform hand hygiene.
- 2. Don non-sterile gloves. Use CHG wipes to clean the tissue around the meatus and outside of the catheter. To avoid contaminating the urinary tract always clean wiping away from, NEVER toward the urinary meatus. Use wipe once and throw it away, cleansing only one area with each wipe.

Do not pull on catheter or the contaminated area will re-enter the urethra.

3.

4. Ensure catheter is properly secured to upper thigh. with securement device.

Anchoring the catheter prevents in and out movement and decreases risk of infection.

5. Teach patient to do above if able.

6. Indwelling foley catheter care and documentation should be done every shift and after bowel movements.

The date and time of foley catheter care should be documented in medical record.

Revised by: Rachel O'neil Cooke, MSN, RN, CVRN-BC, Assistant Unt Director HVCU

Approved: 7/05

Reviewed: 4/08, 8/31/2010, 5/14, 7/14, 2/17, 2/20

Revised: 3/15, 2/20, 10/2024

Reviewed by Policy & Standard Committee, 8/10, 7/14, 2/17, 2/20, 6/23, 10/2024