

## NURSING SERVICES GUIDELINE GENERAL

**Guideline:** Safety Management of the  
Traumatic Brain Injury (TBI)  
or Closed Head Injury (CHI)



**Policy Number Superseded:**

**Effective Date:**  
December 2022

**Responsibility:**

**Initial Effective Date:**  
December 2002

**Purpose of Guidelines:** To protect the patient's rights, to provide for the least restrictive environment, and to provide guidelines for safe and appropriate use of net beds in the treatment of traumatic brain injuries and closed head injuries.

**Equipment:**

- (1) Net bed
- (2) Gait belts
- (3) Wheelchair seatbelts
- (4) Bed alarms
- (5) Other least restrictive devices

**Procedure:** Medical Assessment

- (1) Assessment and correction of underlying medical conditions including iatrogenic causes (especially drugs) which may propagate this acute state of confusion.
- (2) Use of medications to enhance or speed cognitive recovery.
- (3) Appropriate pain management.
- (4) Correction of sleep-wake cycle abnormalities.

**Points of Emphasis:** People with acute brain disorders (TBI, subarachnoid hemorrhage, cerebral tumors, CHI) often undergo a period of acute confusion state/delirium in the process of their recovery. This is a transient medical condition seen in association with post-traumatic amnesia in the trauma setting. Management of patients in this phase of recovery is interdisciplinary involving medical, nursing and therapy services.

## Guideline:

### Safety Management of the Traumatic Brain Injury (TBI) or Closed Head Injury (CHI) 2

PROCEDURE	POINTS OF EMPHASIS
<p>(1) The following baseline data should be assessed by the RN as presenting a safety risk to the patient prior to considering interventions, including restraining devices:</p> <ul style="list-style-type: none"> <li>(a) Fall risk potential.</li> <li>(b) Cognitive functioning.</li> <li>(c) Memory problems.</li> <li>(d) Disorientation/confusion.</li> <li>(e) Impaired judgement.</li> <li>(f) Impulsiveness.</li> <li>(g) Weakness/paralysis.</li> <li>(h) Current medications.</li> <li>(i) Potential or actual fluid and/or electrolyte imbalance.</li> <li>(j) Deformities or medical conditions that would contraindicate the use of restraining devices and/or warrant more careful monitoring.</li> </ul>	<p>Assessment should be done on all patients on admission, and ongoing as needed to prevent patient harm. Refer to the "Fall Assessment" section in daily charting, and Restraints policy <a href="#">3364-100-53-12</a>.</p> <p>Consider any change in patient condition that may warrant physician notification, as cause may require immediate medical treatment.</p> <p>Examples are (but not limited to): Restlessness may be from hypoxia, brain injury, metabolic/endocrine disorders etc. Confusion may be a symptom of hypoxia, hypotension, stroke, TIA, brain injury, hypoglycemia, etc.</p>
<p>(2) Document and inform patient care providers assessment findings of above assessment.</p>	
<p>(3) Consider least restrictive alternative measures and implement as appropriate.</p>	<p>There may be situations when alternative measures are not possible due to the patient condition (i.e., a violent patient). In these situations, documentation as to why least restrictive measures were not attempted must be made in the medical record.</p>
<p>(4) Least restrictive alternative measures to consider:</p> <ul style="list-style-type: none"> <li>(a) Placement of patient close to the nursing station (room or chair).</li> <li>(b) Availability of a sitter/Family Sitter Program.</li> <li>(c) Increased frequency of observation or direct continuous observation 1:1 care.</li> <li>(d) Placement of the call light within the patient's reach.</li> <li>(e) Placement of the side rails in the up</li> </ul>	

## Guideline:

### Safety Management of the Traumatic Brain Injury (TBI) or Closed Head Injury (CHI) 3

<p>position, and the bed in the low position, as needed, based on assessed need.</p> <ul style="list-style-type: none"><li>(f) Consultation with the physician regarding possible medication changes.</li><li>(g) Decreasing sensory stimulation, (i.e. lowering lights, decreasing noise).</li><li>(h) Bed alarms and chair alarms.</li><li>(i) Participation of family in care process.</li><li>(j) Diversional activities such as TV or music.</li></ul>	
<p>(5) Restraining requiring time-limited orders. RN obtains time-limited order for non-violent or violent restraint. Refer to Restraint policy <a href="#">3364-100-53-12</a>.</p>	<p>Net beds utilized for the treatment of a CHI or TBI that present a risk for harm to patient or to others, requires a timed physician order.</p> <ul style="list-style-type: none"><li>▪ When obtaining a time-limited order, include date, time, type or restraints, location and specific reason for the restraint.</li><li>▪ The patient needs to be re-evaluated every calendar day, and only if needed are restraints reordered.</li></ul>
<p>(6) Care of the patient in net bed.</p> <ul style="list-style-type: none"><li>(a) Inform patients and families of the rationale for net bed and necessary changes in patient behavior and/or condition to discontinue. Rationale for the use of the net bed includes:<ul style="list-style-type: none"><li>▪ Patient safety.</li><li>▪ Medications that may potentially impair functioning.</li><li>▪ Behavioral condition.</li><li>▪ Impaired cognitive functioning.</li></ul></li><li>(b) Document in the medical record.<ul style="list-style-type: none"><li>▪ The behavior and/or condition that warrants the need for the net bed.</li><li>▪ The least restrictive interventions utilized.</li><li>▪ A physical assessment of the patient to</li></ul></li></ul>	<p>Clarify with family the patient condition that warrants use of the net bed.</p> <p>Continue to comfort, reassure and offer explanation to patients and families. Families can also be offered the chance to participate in the Family Sitter Program to decrease the time needed for a patient to be in a net bed.</p> <p>Document at least q shift, behavior that warrants continued need for the net bed.</p> <p>Maintain respect and privacy as much as possible.</p> <p>Unsafe items include, but are not</p>

## Guideline:

### Safety Management of the Traumatic Brain Injury (TBI) or Closed Head Injury (CHI) 4

<p>determine the appropriateness of restraints.</p> <p>(c) Nursing care to be provided while the patient is in a net bed and its documentation at least q 2 hours:</p> <ul style="list-style-type: none"><li>▪ Respiratory/circulation status.</li><li>▪ Emotional and supportive care provided.</li><li>▪ Fluid and foods offered.</li><li>▪ Toileting offered at least every two hours.</li><li>▪ Skin care/attention given to skin, turning repositioning, offer bathing every 24 hours and more frequent hygiene as needed.</li><li>▪ Elevate head of bed to reduce the possibility of aspiration.</li><li>▪ Remove unsafe objects from patient's reach.</li></ul>	<p>limited to: smoking materials, sharp objects, medications and other potentially dangerous materials.</p> <p>Any redness bruising or changes in patient condition need to be reported to the physician and documented.</p>
<p>(7) Discontinuation of the net bed.</p> <p>(a) A net bed will be discontinued when the patient no longer presents a risk for harm to self or others. This would mean prior to the expiration.</p> <p>(b) If assessment warrants early release from the net bed, rationale for discontinuation could include:</p> <ul style="list-style-type: none"><li>▪ Availability of sitter.</li><li>▪ Behavioral changes.</li><li>▪ Least restrictive measures that were successful.</li></ul> <p>(c) When discontinuing the net bed:</p> <ul style="list-style-type: none"><li>▪ Note time discontinued.</li><li>▪ Note behavior of patient.</li><li>▪ Rationale for discontinuing.</li><li>▪ Status of the patient following removal up to the first sixty minutes.</li></ul> <p>(d) When the net bed is terminated early, and the same reason for the net bed is</p>	<p>Because restraints can affect the dignity of patients, as well as causing harm, the decision to restrain or not should be discussed during team conference.</p> <p>Document: Date and time of every episode of application or discontinuation of restraints.</p>

## Guideline:

Safety Management of the Traumatic Brain Injury (TBI) or Closed Head Injury (CHI) 5

still evident, a new order must be obtained if alternatives are ineffective.	
--	--

References: 2022 CMS Standards  
Joint Commission Provision of Care Standards 2022

Approved by:  
*Kurt Kless, MSN, MBA, RN, NE-BC*  
*Chief Nursing Officer*

Initial effective date:  
December 2002

*Review/Revision Completed by:*  
*Mike Drake, BSN, RN*  
Reviewed by Policy & Standard  
Committee  
*August 2012, July 2016, September*  
*2019, December 2022*

Review/Revision Date:  
June 2005  
September 2008  
August 2012  
July 2016  
September 2019  
December 2022

Next review date:  
December 2025