

NURSING SERVICE GUIDELINES INPATIENT BEHAVIORAL HEALTH

Guideline: Psychotic patient

Policy Number Superseded:

Responsibility:

Purpose of Guideline:



Effective Date:

June 2023

Initial Effective Date:

September 1993

Psychosis describes a degree of severity not a specific disorder. A psychotic patient has a grossly impaired sense of reality often coupled with emotional and cognitive disabilities, which severely compromises his ability to function. He is likely to talk and act in a bizarre fashion, hallucinate or strongly hold ideas that are contrary to fact (delusions.) He may be confused and disoriented.

PROCEDURE	POINT OF EMPHASIS
(1) Assess for symptoms of psychosis on admission or during hospitalization: (a) Delusions. (b) Hallucinations. (c) Disorganized thinking or speech. (d) Flat emotional affect. (e) Lack of goal directed activity. (f) Limited productive thoughts.	
(2) Patient can expect care directed toward preventing complications or injury of the psychotic patient. Patient can expect to be helped to maintain contact with reality as much as possible.	(2) The nurse will: (a) Use medication as ordered. Low dose antipsychotic drug maintenance is the norm. (b) Be specific. Ask pointed, factual questions. Try to identify the patient's major current fears and concerns, but don't be led into a lengthy discussion of complex delusions and hallucinations. Use

	<p>simple words and state that patient may hear voices, but you don't.</p> <p>(c) Don't rush the patient to respond to each question but do maintain some control over the direction of the conversation.</p> <p>(d) Make some specific observations of the patient's behavior (e.g., "you look frightened, you look angry") but don't become involved in lengthy interpretations. Don't draw incorrect conclusions about the emotional state from inappropriate effect.</p> <p>(e) Explain to the patient what is being done to him and why.</p> <p>(f) If the conversation is going nowhere, e.g., the patient refuses to talk, break off the interview with a positive expectation, e.g., "I'll be back to see you in a little while when you are feeling better and able to talk."</p> <p>(g) Keep environment quiet.</p>
(3) Avoid touching the patient especially a paranoid patient.	
(4) Keep the unit structured with concrete activities.	
(5) Teach coping skills: (a) Including assertiveness. (b) Participation with others. (c) Anxiety relief activities. (d) Deep breathing.	
(6) Document clearly any hallucinations, patient thoughts and actions.	
(7) Observe for suicidal behavior.	
(8) If patient becomes aggressive attempt least restrictive interventions.	(8) Use of restraints when needed: Restraints are used only when less restrictive techniques have been

	exhausted and when patient continues to exhibit behavior injurious to himself or others. Refer to restraint/seclusion guideline.
--	--

(A) References.

Van Schalkwyk, G. I., Davidson, L. & Srihari, V. (2015). Too late and too little: Narratives of treatment disconnect in early psychosis. *Psychiatric Quarterly*, 86(4), 521-532.

Ajnakina, O., Trotta, A., Oakley-Hannibal, E., Di Forti, M., Stilo, S. A., Kolliakou, A., et al. (2016). Impact of childhood adversities on specific symptom dimensions in first-episode psychosis. *Psychological Medicine*, 46(2), 317-326.

Stone, J. M., Fisher, H. L., Major, B., Chisholm, B., Wooley, J., Lawrence, J., et al. (2014). Cannabis use and first-episode psychosis: Relationship with manic and psychotic symptoms, and with age at presentation. *Psychological Medicine*, 44(3), 499-506.

Mosby's Nursing Skills 2020

Approved by:
Kurt Kless, MSN, MBA, RN, NE-BC
Chief Nursing Officer

Initial effective date:
September 1993

Review/Revision Completed by:
Tamara Cerrone, BSN, RN, Director
Inpatient Behavioral Health
Reviewed by Policy & Standard
Committee
August 2011, August 2014, July 2017,
July 2020, June 2023

Review/Revision Date:
May 1996
April 1999
March 2002
April 2005
June 2008
August 2011
July 2017
July 2020
June 2023

Next review date:
June 2026