

Nursing Service Guidelines Inpatient Behavioral Health



Title: CARE OF THE PATIENT WHO HAS AN ANXIETY DISORDER

Description: Anxiety is a diffuse, highly unpleasant, often vague feeling of apprehension, accompanied by one or more bodily sensations; for example, pounding heart, perspiration, headache or the sudden urge to void. Anxiety is an alerting signal; it warns of impending danger and enables the person to take measures to deal with a threat. Anxiety is in response to a threat that is unknown, internal, vague, or conflicting in origin.

Purpose:

1. The patient will understand, accept, and learn to live with anxiety.
2. The patient will learn methods and skills to cope constructively with anxiety.

Procedure

Point of Emphasis

- | | |
|---|---|
| 1. Assess level of anxiety. Observe for behavior characteristics of anxiety. Restlessness, easily fatigued, problems concentrating, irritability, tension and sleep problems. Anxiety is a normal response to situations, however, becomes a problem if it interferes with adaptive behavior, causes physical symptoms. | To identify anxiety. Changes in vital signs could occur such as blood pressure increases, increase in pulse, respirations, tightening of throat, epigastric pain. Work with the patient on identifying issues early in hospitalization so that treatment interventions can begin. |
| 2. Ask, “what are you feeling?” | To help the patient name the feeling. |
| 3. Connect the feeling with behavior. Patient will attend scheduled groups and address the following issues in those groups: <ul style="list-style-type: none">• Explore with the patient what happened prior to her feeling anxious.• Discuss alternatives for dealing with the situation (cause). Assessments, nursing care, and patient teaching must be documented on the daily assessment checklist and nursing progress notes. Assist patient in identifying precipitating factors to reduce frequency and intensity of anxiety. | To discover the cause.

To improve patterns of handling anxiety. |
| 4. Reduce stimuli as it can create confusion, also increasing anxiety. Speak in short, clear sentences. | As anxiety progresses it results in decreased concentration and difficulty in decision making and problem solving. |
| 5. Teach alternatives to handling anxious situations such things include: <ul style="list-style-type: none">• Breathing techniques• Imaging• Visualization | Help patient gain control over situation and decrease anxiety. |
| 6. Teach/educate parent(s) and the child about the anxiety disorder. | |
| 7. Educate parent(s) and the child regarding outpatient treatment. | |

Care of the Patient Who Has an Anxiety Disorder Guidelines
Page 3

Reviewed by: Tammy Cerrone, BSN, RN, Nursing Director Kobacker

Approved: September 1993

Reviewed: 4/99, 3/02, 9/17, 9/2020, 10/2023

Revised: 4/96, 5/05, 6/08, 6/10, 8/14

Reviewed by Policy & Standard Committee: 3/10; 8/14, 9/17, 9/2020, 10/2023

References: Manual of Psychiatric Nursing Care Plans Diagnoses, Clinical tools and Psychopharmacology, 2nd Edition, Varcavolis, Elizabeth M. 2013.
www.guidelines.gov/summary ; Connolly, S.D., Berstein, G.A., Work Group on Quality Issues. Practice parameter for the assessment and treatment of children and adolescents with anxiety disorders. J Am Acad Child Adolesc Psychiatry 2007 Feb; 46(2): 267-83. (192 References)
Pediatric Nursing, July-August 1992, Vol. 18, No. 4. The 'acting out' adolescent: identification and management.
Connolly, S.D., Nanayaleara, S.D. 2009. "Anxiety Disorders in Children and Adolescents: Early Identification and Evidence-Based Treatment". *Psychiatric News*